



Schweizerische Arbeitsgemeinschaft für Klinische Krebsforschung
Groupe Suisse de Recherche Clinique sur le Cancer
Swiss Group for Clinical Cancer Research
Gruppo Svizzero di Ricerca Clinica sul Cancro



Rapporto annuale

La versione PDF del rapporto annuale 2008 è disponibile su
<http://sakk.ch>

Indice

Editoriale	
Attualità 2008	2
Gruppo Svizzero di Ricerca Clinica sul Cancro	
Attività del centro di coordinamento	4
Membri del consiglio direttivo	7
Organigramma	8
Forze dirigenziali, onorificenze e promozioni all'interno del SAKK	9
SAKK Assemblea semestrale 2008	10
Scientific Activities	
Summary of Activities, Trials	12
Project Group Reports	
Project Group Breast Cancer	15
Project Group Gastrointestinal Cancer	16
Project Group Leukemia	19
Project Group Lung Cancer	21
Project Group Lymphoma	22
Project Group New Anticancer Drugs/Phase I Trials	24
Project Group Urogenital Tumors	25
Sections	
Section Clinical Research Coordinators (CRC)	26
Section Internal Medicine	27
Section Pathology	27
Section Radio-Oncology	28
Section Surgery	30
Section Trial Nurses	31
Networks	
Network for Cancer Predisposition Testing and Counseling (CPTC)	32
Network for Outcomes Research	33
Publications 2008	35
Resoconti annuali	39
Indirizzi	40



Attualità 2008

del Prof. Richard Herrmann | Presidente del SAKK

Il 2008 è stato un buon anno per il SAKK. Siamo riusciti a raggiungere diversi obiettivi tra quelli che ci eravamo posti. E il rapporto annuale parla proprio di questo.

Il 2008 è stato il primo anno in cui il SAKK ha riguadagnato la completa autonomia. E gli effetti positivi non si sono visti solamente nel fatto che i suoi soci hanno avuto bisogno di incontrarsi meno di prima. Si percepisce anche nella considerazione guadagnata dal lavoro svolto dal SAKK, tanto dai suoi membri quanto dal Centro di Coordinamento, e dall'importanza che è riconosciuta alla nostra attività.

Quanto sia difficile organizzare una ricerca clinica lo stanno vedendo ora le Clinical Trial Units (CTU) di recente fondazione. Queste unità, sostanzialmente, agiscono ancora a livello locale. Il fatto che noi siamo parecchio più avanti in questo campo lo dobbiamo alla lungimiranza della generazione fondatrice del SAKK.

Qual è il «segreto» che ci permette di funzionare così bene? È l'interazione di diverse forze:

- da un lato, i membri di tutte le regioni della Svizzera e dei membri associati esteri,
- dall'altro lato, il Centro di Coordinamento del SAKK di Berna.

Perché queste forze possano interagire, abbiamo bisogno:

- di obiettivi comuni,
- di comprensione per il modo di operare degli altri e per i loro problemi e
- di una comunicazione aperta.

L'auspicio di tutti i responsabili del SAKK è quello di trovare un giusto equilibrio fra tutti gli interessi in campo. Dobbiamo soddisfare le richieste dei nostri committenti e le esigenze dei nostri membri, ma anche quelle del nostro stesso Centro di Coordinamento, il cui carico di lavoro deve rimanere in un ambito gestibile, ma sufficiente a consentirci di raggiungere i nostri obiettivi. Vorremmo portare avanti in fretta i progetti già approvati, mantenendo però la qualità del lavoro e della vita – fattori a cui teniamo moltissimo – delle nostre collaboratrici e dei nostri collaboratori.

È incoraggiante il fatto che l'anno scorso in alcuni *project group* siano state affidate responsabilità di elaborazione di *concept* e realizzazione di studi clinici a numerosi giovani colleghe e colleghi. Questa tendenza dovrebbe mantenersi. Pertanto invito questi giovani a discutere nuove idee nell'ambito dei loro gruppi e a lanciare nuovi progetti. Per frequentare corsi speciali sulla ricerca clinica e per brevi soggiorni presso centri di ricerca all'estero ci sono ancora delle risorse a disposizione.

Affinché si possa continuare a fare ricerca clinica d'eccellenza, il Consiglio direttivo ha proposto ai gruppi di progetto di invitare, come interlocutori, uno o più esperti internazionali del campo. Riteniamo che, in questo modo, nei nostri progetti possano entrare – più in fretta e meglio supportate – idee nuove.

L'anno scorso abbiamo avuto diversi colloqui, molto costruttivi, con autorità e organizzazioni che ci sostengono (Segreteria di Stato per l'educazione e la ricerca) o con le quali lavoriamo per l'ottimizzazione della ricerca clinica in Svizzera (Swissmedic, Comunità di lavoro delle commissioni etiche cantonali della Svizzera [AGEK]). Sono certo che sapremo risolvere insieme i problemi che ci attendono.

I nostri obiettivi per questo 2009 rimangono invariati: vogliamo effettuare studi clinici di eccellenza in campo oncologico che siano competitivi a livello internazionale. Vogliamo fare in modo che i pazienti oncologici della Svizzera abbiano accesso a nuove terapie capaci di dar loro nuove speranze; vogliamo anche dare il nostro contributo, a livello internazionale, al miglioramento dei risultati delle cure.

Mi auguro che tutti gli interessati collaborino per il raggiungimento di questi traguardi in modo fattivo e impegnato come hanno fatto l'anno scorso. Ne vale la pena. Siamo riusciti a pubblicare i risultati del nostro lavoro su riviste specializzate di importanza internazionale, quali il *Journal of Clinical Oncology* – dove siamo stati presenti più di una volta – e il *Lancet Oncology*. Sono apparse in tutto 44 pubblicazioni su studi portati a termine dal SAKK o studi ai quali il SAKK ha partecipato in modo sostanziale. Questo rispecchia l'efficienza del SAKK e la sua influenza nel campo delle ricerche oncologiche.

Il 2008 è stato per noi un anno proficuo e questo lo dobbiamo all'impegno di tutti i membri, di tutte le collaboratrici e tutti i collaboratori del Centro di Coordinamento e ad ogni singolo membro delle istituzioni che aderiscono al SAKK. Colgo l'occasione per esprimere a tutti loro i miei personali ringraziamenti.

4 | Gruppo Svizzero di Ricerca Clinica sul Cancro SAKK



Attività del Centro di Coordinamento

del Dott. Peter Brauchli | Direttore del SAKK

Dopo una ristrutturazione, spesso ci si chiede se ne valeva la pena. A questa domanda possiamo senz'altro rispondere «sì». I doppioni che c'erano prima causavano perdite di tempo; la fusione avvenuta successivamente fra SIAK e SAKK ha richiesto, negli anni passati, l'impiego di molte risorse. Ora abbiamo di nuovo più tempo per lavorare in modo produttivo. È possibile e necessario, comunque, apportare ulteriori miglioramenti nella nostra attività quotidiana, in modo da raggiungere più in fretta i traguardi che ci siamo dati. A tale proposito è importante ricordare il nostro obiettivo principale: compito del SAKK è elaborare e attuare studi clinici d'eccellenza in ambito oncologico, i cui risultati possano essere messi a disposizione dei pazienti nel minor tempo possibile.

Alcuni compiti importanti sono già in fase di realizzazione, mentre alcuni dei progetti per il futuro sono in fase di ulteriore sviluppo. Tra questi ci sono l'elaborazione delle Standard Operating Procedures (SOPs), la realizzazione del dipartimento di Regulatory Affairs e l'ampliamento del nostro Electronic Data Capturing System SINATRAS. Questi cambiamenti, adattamenti e miglioramenti sono necessari per avere successo in futuro. Inoltre, per il SAKK è indispensabile posizionarsi in modo chiaro e far valere i propri diritti nel campo della ricerca clinica in Svizzera, dove i cambiamenti si susseguono rapidamente. Perciò, nel corso dell'anno abbiamo definito chiaramente che cos'è il SAKK

e quale è la sua attività, in modo che quando se ne parla, si abbia un'idea chiara di questa realtà. Speriamo, in questo modo, di riuscire a potenziare anche nei vari centri la consapevolezza del significato della parola «noi».

Il posizionamento del SAKK

Cosa facciamo e chi siamo

Il SAKK è un'associazione senza scopo di lucro, che si dedica dal 1965, come gruppo cooperativo, alla ricerca clinica oncologica sul territorio nazionale.

Sulla base di un contratto di prestazione stipulato con la Confederazione, il SAKK ha funzione di istituto di ricerca universitario decentralizzato, effettua studi clinici nei maggiori ospedali della Svizzera e collabora, a livello internazionale, con gruppi cooperativi stranieri.

Il suo scopo è quello di sviluppare ulteriormente le terapie oncologiche in pazienti adulti – uomini e donne – e studiare l'efficacia e la tollerabilità di nuove terapie.

Il SAKK, con il suo network, è interlocutore privilegiato di istituzioni, associazioni, aziende e case farmaceutiche svizzere nel campo della ricerca clinica oncologica.

Nella lotta contro il cancro condotta a livello nazionale, il SAKK collabora con tutti questi organismi.

Consiglio direttivo del SAKK

Il Consiglio direttivo del SAKK, rispetto all'ex Comitato Esecutivo, ha maggiori responsabilità e nuovi compiti, che svolge nel rispetto dei principi della corporate governance delle istituzioni non a scopo di lucro. Affinché la scelta degli studi da parte del SAKK possa avvenire in conformità ai criteri internazionalmente riconosciuti, i gruppi di progetto e gli altri organi specializzati hanno il diritto di presentare le loro istanze.

Per quanto riguarda la scelta degli studi, nel corso dell'anno il Consiglio direttivo ha perfezionato il proprio modo di procedere. Esso decide in merito alla possibilità di effettuare un determinato studio in seduta plenaria, attenendosi ai seguenti criteri:

- a) rilevanza scientifica,
- b) fattibilità (inclusa la stima di reclutamento),
- c) conformità alla strategia del SAKK,
- d) finanziabilità.

Questa scelta diventa tanto più difficile, quanto più numerosi sono i responsabili di studi che elaborano progetti di grande validità e li presentano al Consiglio direttivo. Quattro studi, che nel 2008 avevano ricevuto una valutazione definitiva, non sono poi stati realizzati. Le ragioni della mancata attuazione sono state, principalmente, il venir meno del consenso del Consiglio direttivo del SAKK e la carenza del supporto fornito dalle aziende interessate. Per mettere ordine nelle procedure di scelta, sempre più complesse, è stato elaborato un regolamento organizzativo.

Inoltre, ad alcuni membri del Consiglio direttivo è stata affidata la responsabilità di una o più sezioni. I contatti più importanti fra il Consiglio direttivo e i gruppi di progetto vengono gestiti attraverso un cosiddetto referente (liaison person), il cui compito essenziale è quello di rappresentare l'opinione del Consiglio direttivo nei gruppi di progetto e di trasmettere le raccomandazioni del Comitato scientifico ai gruppi di progetto tramite il Consiglio direttivo. Così anche i consulenti internazionali e gli esperti dovrebbero partecipare più numerosi alle riunioni dei gruppi di progetto, mentre i membri dovrebbero sentirsi incoraggiati a presentare ai congressi gli *abstract* degli studi del SAKK.

Strategia

In aprile il Consiglio direttivo del SAKK ha ridefinito, durante un ritiro appositamente organizzato, l'orientamento strategico e sviluppato le raccomandazioni del Comitato scientifico. Il gruppo di progetto Tumori Uro-Genitali fondato su tali raccomandazioni ha potuto avviare, nel 2008, un primo studio, a cui faranno seguito almeno altri due nel 2009. Durante il ritiro sopraccitato si è anche discusso del

fatto che i medici che effettuano gli studi dovrebbero venire incoraggiati ad occuparsi di più degli studi di fase III relativi alle indicazioni più importanti, includendo anche aspetti di ricerca traslazionale (translational research), di impatto dei risultati (outcomes research), di qualità della vita. Diversi studi promettenti di fase II dovrebbero aprirci la possibilità di valutare in seguito le terapie testate che hanno dato risultati positivi anche in studi di fase III.

Collaborazione con centri stranieri

La collaborazione con centri stranieri si rivela spesso difficile. Le cause sono molteplici. Aprire un centro all'estero è costoso e comporta ritardi nella partenza degli studi, spesso perché l'azienda farmaceutica non metterà con certezza a disposizione il medicinale oggetto di studio nel Paese in cui il centro dovrebbe essere attivato. Per meglio affrontare questo problema è stato creato e reso operativo, presso il Centro di Coordinamento del SAKK, l'ufficio «Regulatory Affairs». Tra tutti i centri stranieri, l'IEO di Milano continua ad essere il nostro partner più importante. La collaborazione prosegue positivamente anche con il Centro di Budapest. Friburgo in Breisgau (D) ha incluso un primo paziente in uno studio SAKK. Nel corso del 2009 ci saranno incontri con rappresentanti dei centri esteri, finalizzati a far includere a questi ultimi più pazienti nei loro studi.

L'attività di studio

L'anno scorso il SAKK ha incluso 817 pazienti in 44 studi clinici, ovvero il numero più alto degli ultimi cinque anni. Ben 532 pazienti sono stati reclutati in studi nostri e 285 in studi di gruppi cooperativi stranieri. Nel 2008 abbiamo gestito complessivamente un numero di nuovi protocolli che è il più alto di sempre. Il Consiglio direttivo ha accettato, complessivamente, sei studi nostri e cinque studi di gruppi cooperativi esteri. Nel 2008 il SAKK ha avviato cinque nuovi studi.

L'anno scorso sono stati pubblicati 44 lavori, dei quali il SAKK era responsabile o ai quali aveva partecipato in modo determinante. Questo è il numero più alto negli oltre

40 anni di storia del SAKK. Inoltre, siamo orgogliosi del fatto che il SAKK ha partecipato a due studi, i cui risultati sono stati giudicati da una commissione indipendente dell'American Society of Oncology (ASCO) come significativo passo in avanti e importanti per scopi terapeutici (terapia adiuvante del cancro al seno (NCIC CTG MA.17. IBCSG Trial 17-98) e come terapia adiuvante del melanoma (EORTC 18991)).

Cooperazioni

Con la *Clinical Trial Unit* (CTU) di Berna il SAKK ha attuato, nel 2008, il primo aggiornamento per medici responsabili di studi. Inoltre, il dipartimento statistico del SAKK ha organizzato, insieme all'Istituto di medicina sociale e preventiva (ISPM) dell'Università di Zurigo, un simposio dal titolo «Standardizzazione e individualizzazione». Entrambi gli eventi hanno suscitato grande interesse, ed è stata buona la partecipazione di diversi gruppi professionali.

È stato anche elaborato un progetto di cooperazione con lo Swiss Institute of Bioinformatics (SIB) di Losanna per l'analisi dei dati relativi ai marcatori tumorali. Abbiamo avuto incontri molto proficui con la Swiss Clinical Trial Organization (Swiss-CTO).

Il GOPS ha rafforzato le proprie strutture con la creazione di una propria segreteria presso il Centro di Coordinamento del SAKK e la nomina di una nuova direttrice, e ora può operare ancora meglio come organizzazione autonoma. Allo stesso tempo il GOPS continua a collaborare con il SAKK soprattutto nell'ambito delle relazioni con le istituzioni esterne.

Centro di Coordinamento

Per far fronte al crescente numero di studi da sviluppare e, di conseguenza, al lavoro che questi comportano, alla fine del 2008 il numero degli impiegati in pianta stabile del SAKK è stato aumentato di otto unità ed ha raggiunto quota 41 (3395 punti percentuale)

Il SAKK ha assunto diverse persone e affittato nuovi locali soprattutto per seguire il campo principale di attività (sviluppo e attuazione di studi). Oltre al coordinamento degli studi, sono stati potenziati anche i dipartimenti di Statistica e Informatica. La Direzione del Dipartimento «Regulatory Affairs» è stata affidata a Pascale Wenger, mentre Edith Rickenbacher è la nuova responsabile del Dipartimento Amministrazione, Finanze e Personale.

Membri del consiglio direttivo



Presidente

Prof. Richard Herrmann
Universitätsspital Basel



Vizepresidente

Prof. Beat Thürlimann
Kantonsspital St. Gallen



Prof. Daniel Betticher
Kantonsspital Freiburg



Prof. Stephan Bodis
Kantonsspital Aarau



PD Dott. Yves Chalandon
Hôpital Universitaire
de Genève



Prof. Martin Fey
Inselspital Bern



Prof. Michele Ghielmini
Ospedale Regionale
Lugano



Prof. Holger Moch
Universitätsspital Zürich



Prof. Christoph Renner
Universitätsspital Zürich



PD Dott. Arnaud Roth
Hôpital Cantonal
Universitaire Genève




Dott. Roger von Moos
Kantonsspital Chur



Prof. Walter Richard Marti
Universitätskliniken Basel

Organigram Swiss Group for Clinical Cancer Research (SAKK)



Forze dirigenziali, onorificenze e promozioni all'interno del SAKK

Primari

- Prof. Thürlimann, primario del Centro Tumori del Seno, Ospedale Cantonale di San Gallo
- Libero Docente Dott. Abdelkarim S. Allal, primario di Radio-Oncologia, Ospedale Cantonale di Friburgo
- Libero Docente Dott. Emanuele Zucca, vice primario del Dipartimento di Ricerca, IOSI, Istituto di oncologia della Svizzera italiana, Ospedale San Giovanni, Bellinzona

Dirigenti medici

- Dott. Thomas Ruhstaller, dirigente medico del Centro Tumori del Seno, Ospedale Cantonale di San Gallo: dall' 1.1. 2009
- Prof. Jens Huober, capoclinica con funzioni straordinarie, Centro Tumori del Seno, Ospedale Cantonale di San Gallo: dall' 1.1. 2009)
- Prof. Dott. Bernhard Pestalozzi, dirigente medico a. i., Clinica e Policlinico Oncologico, Clinica Universitaria di Zurigo

Libero Docente

- Dott. Florian Strasser, capo clinica di Oncologia e Medicina Palliativa, Dipartimento Medicina Interna, Ospedale Cantonale di San Gallo
- Dott. Viviane Hess, clinica di Oncologia, Clinica Universitaria di Basilea

Cattedra universitaria

- Libero Docente Dott. Bernhard Pestalozzi, professore titolare, Clinica e Policlinico Oncologico, Clinica Universitaria di Zurigo
- Libero Docente Dott. Christoph Driessen, dirigente del Dipartimento di Ricerca Clinica, Ospedale Cantonale di San Gallo
- Libero Docente Dott. Martin H. Brutsche, primario Pneumologia, Ospedale Cantonale di San Gallo

Consiglio direttivo

- Prof. Dott. Walter Richard Marti, dirigente medico, Clinica Universitaria di Basilea
- Prof. Dott. Michele Ghielmini, Istituto oncologico della Svizzera italiana, Ospedale San Giovanni, Bellinzona

Gruppi di progetto SAKK

- Prof. Christoph Rochlitz, Dipartimento di Oncologia, Clinica Universitaria di Basilea: co-presidente del gruppo di progetto sul tumore al seno (successore del Prof. Stefan Aebi)

Sezioni del SAKK

- Prof. Paul M. Schneider, Clinica di Chirurgia Viscerale e dei Trapianti, Clinica Universitaria di Zurigo: presidente della Sezione Chirurgia del SAKK (successore del Prof. Walter R. Marti)
- Prof. Bernhard Pestalozzi, Clinica e Policlinico Oncologico, Clinica Universitaria di Zurigo: Presidente del Network for Outcomes Research
- Prof. Thomas Szucs, Istituto di Medicina Sociale e Preventiva, Università di Zurigo: Vicepresidente Network for Outcomes Research

Nomine

- Libero Docente Dott. Markus Manz, vicedirettore dell'IRB (Istituto di ricerca in Biomedicina), Bellinzona

Premi

- Prof. Franco Cavalli, Direttore dell'Istituto Oncologico della Svizzera italiana, Ospedale San Giovanni, Bellinzona: Premio della Lega Svizzera contro il cancro
- Libero Docente Dott. Emanuele Zucca, vice primario del Dipartimento di Ricerca, IOSI, Istituto oncologico della Svizzera italiana, Ospedale San Giovanni, Bellinzona: Premio SAKK-Pfizer
- Dott. Roger von Moos, Oncologia Medica, Ospedale cantonale di Coira: Premio SAKK-Pfizer

Assemblea semestrale di giugno

L'assemblea del SAKK per il semestre estivo si è tenuta il 19 giugno presso il Centro Congressi Blumenberg di Berna. Hanno partecipato all'incontro circa 150 oncologi, 50 rappresentanti delle case farmaceutiche e 30 collaboratori del Centro di Coordinamento del SAKK. Le riunioni dei gruppi di progetto sono state molto frequentate. Anche i gruppi di lavoro e le sezioni si sono incontrati nell'ambito dell'assemblea.

Il network Outcomes Research ha eletto un nuovo presidente, Bernhard Pestalozzi, e un nuovo vicepresidente, Thomas Szucs.

Il Prof. Walter Richard Marti della Clinica Universitaria di Basilea è stato nominato nuovo membro del Consiglio direttivo dall'Assemblea dei membri del SAKK in sostituzione di Prof. Markus Zuber dell'Ospedale Cantonale di Olten, ritiratosi dopo due mandati.

Assemblea semestrale di novembre

Il 20 e il 21 novembre si sono incontrati a Basilea per l'assemblea semestrale invernale quasi 200 fra oncologi, collaboratrici del SAKK e rappresentanti dell'industria farmaceutica. Durante il convegno si sono tenuti due simposi con interventi interessanti. Le sessioni dei gruppi di progetto del SAKK, dei gruppi di lavoro e delle sezioni, nei quali sono stati discussi i progetti di ricerca in essere e quelli futuri, hanno riscosso notevole interesse.

Al simposio del SAKK è intervenuto il Prof. Ruggero De Maria dell'Istituto Superiore di Sanità di Roma. Ha parlato delle cellule staminali e del loro uso mirato nelle terapie anticancro. Poi c'è stata la consegna del Premio SAKK/Pfizer 2008, ammontante a 20.000 franchi svizzeri. Il riconoscimento ai progetti che migliorano la qualità della ricerca clinica in Svizzera è stato assegnato dal Presidente del SAKK, Prof. Richard Herrmann, e dal Dott. Thomas Schaller, in rappresentanza della Pfizer, al Dott. Roger von Moos e al Libero Docente Dott. Emanuele Zucca.

Il Dott. Zucca, primario del Reparto Linfomi dello IOSI, ha ricevuto la sua parte di premio per aver istituito una banca dati per i linfomi. Il Dott. von Moos, dirigente medico di Oncologia e Ematologia dell'Ospedale Cantonale dei Grigioni, è stato premiato per aver avviato la ricerca clinica oncologica nello stesso Canton Grigioni.



Prof. Bernhard Pestalozzi



Prof. Thomas Szucs



Prof. Walter Richard Marti



Prof. Michele Ghielmini



I membri del SAKK hanno acconsentito, in sede di assemblea, a portare i membri del Consiglio direttivo da 11 a 12. È stato eletto all'unanimità, quale rappresentante della Svizzera italiana nel Consiglio direttivo, il Prof. Michele Ghielmini, primario di Oncologia medica presso l'Istituto oncologico della Svizzera italiana (IOSI).



Dott. Thomas Schaller, PD Dott. Emanuele Zucca,
Dott. Roger von Moos, Prof. Richard Herrmann (da sx)

Conferimento del SAKK/Amgen Research Grant 2009

Il premio – ammontante a 50.000 franchi svizzeri – per la ricerca traslazionale e la ricerca oncologica innovativa e orientata al miglioramento della vita dei pazienti verrà conferito a Basilea durante l'assemblea semestrale del SAKK (26/27 novembre).

Il regolamento di partecipazione al SAKK/Amgen Research Grant 2009 può essere richiesto a:

Prof. Richard Herrmann, Presidente SAKK
SAKK Koordinationszentrum
Effingerstrasse 40, 3008 Berna

By Dr Stefanie Lerch | Head of Trial Coordination

Summary of Activities

In 2008, a total of 817 patients (632 in 2007) were included in 44 clinical trials coordinated by SAKK*:

	2008	2007
Total patients from Switzerland	773	584
Total patients from foreign countries	44	48
Total	817	632

	2008	No. of trials
Total patients in SAKK trials	532	21
Total patients in trials of cooperative groups (without IBCSG)	139	13
Total patients in IBCSG trials	100	6
Total patients in Sando trials	46	4
Total	817	44

* SAKK 21/07 and SAKK 63/03 are not included

Trials open for accrual in 2008

Urogenital Cancer

SAKK 08/07 | Docetaxel and cetuximab in patients with docetaxel-resistant hormone-refractory prostate cancer (HRPC). A multicenter phase II trial

Lung Cancer

SAKK 16/00 | Preoperative radiochemotherapy vs. chemotherapy alone in non-small cell lung cancer patients with mediastinal lymph node metastases (stage IIIA, N2). A randomized phase III trial

SAKK 17/04 | Neoadjuvant chemotherapy and extrapleural pneumonectomy of malignant pleural mesothelioma (MPM) with or without hemithoracic radiotherapy. A randomized multicenter phase II trial

SAKK 19/05 | Bevacizumab and erlotinib first-line therapy in advanced non-squamous non-small cell lung cancer (stage IIIB/IV) followed by platinum-based chemotherapy at disease progression. A multicenter phase II trial

Breast Cancer

SAKK 22/99 | Randomized phase III trial of Herceptin® followed by chemotherapy plus Herceptin® versus the combination of Herceptin® and chemotherapy as palliative treatment in patients with HER2-overexpressing advanced/metastatic breast cancer

SAKK 23/03 | Trastuzumab monotherapy followed by the combination of trastuzumab and letrozole in postmenopausal women with ER-positive, HER-2 positive advanced breast cancer resistant to a nonsteroidal aromatase inhibitor. A multicenter two-step phase II trial

SAKK 24/06 | Bevacizumab (Avastin®) and pegylated liposomal doxorubicin (Caelyx®) as first-line therapy for locally recurrent or metastatic breast cancer. A multicenter, open-label phase II trial

IBCSG 22-00 | Low-dose Cytotoxics as «Anti-angiogenesis Treatment» following Adjuvant Induction Chemotherapy for Patients with ER-negative and PgR-negative Breast Cancer

IBCSG 23-01 | A randomized trial of axillary dissection vs. no axillary dissection for patients with clinically node negative breast cancer and micro-metastases in the sentinel node

IBCSG 24-02 | BIG 2-02/SOFT Suppression of Ovarian Function Trial (SOFT). A Phase III Trial Evaluating the Role of Ovarian Function Suppression and the Role of Exemestane as Adjuvant Therapies for Premenopausal Women with Endocrine Responsive Breast Cancer

IBCSG 27-02 | BIG 1-02/NSABP Trial B-37 A randomized clinical trial of adjuvant chemotherapy for radically resected loco-regional relapse of breast cancer

IBCSG 35-07 | BIG 1-07 SOLE Study of Letrozole Extension. A phase III trial evaluating the role of continuous letrozole versus intermittent letrozole following 4 to 6 years of prior adjuvant endocrine therapy for postmenopausal women with hormone-receptor positive, node positive early stage breast cancer

IBCSG 36-07 | ALTTO (Adjuvant Lapatinib and/or Trastuzumab Treatment Optimisation) study. A randomised, multi-centre, open-label, phase III study of adjuvant, lapatinib, trastuzumab their sequence and their combination in patients with HER2/ErbB2 positive primary breast cancer

IBIS II | International Breast Cancer Intervention Study. A randomised double blind control trial divided into two strata

Leukemia

SAKK 30/00/HOVON42 | Randomized induction concerning the value of G-CSF priming in adult patients (≤ 60 yrs of age) with acute myelocytic leukemia (AML) or refractory anemia with excess of blasts (RAEB, RAEB-t) with IPSS score ≥ 1.5

SAKK 30/07 | 5-Azacytidine to treat acute myeloid leukemia in elderly or frail patients not suitable for intensive chemotherapy. A multicenter phase II trial

APL 2006 | A randomized trial assessing the role of arsenic trioxide and/or ATRA during consolidation course in newly diagnosed acute promyelocytic leukemia (APL)

CLL7 | Randomized phase III trial comparing early treatment with fludarabine, cyclophosphamide+rituximab versus deferred treatment in untreated Binet stage A patients with high risk of progression

CML IV | Randomized controlled comparison of Imatinib vs Imatinib/IFN- α vs Imatinib high-dose (800 mg) and determination of the role of allografting in newly diagnosed CML

GRAALL 2005 | Randomized phase III trial assessing the value of intensive vs standard induction and intensification in a randomized comparison and for B-ALL in a second randomization the benefit of rituximab in addition to chemotherapy and for Ph⁺ ALL in a randomized comparison the non-inferiority of an imatinib based induction therapy vs a chemotherapy based induction combined with imatinib

HOVON 81 | A Phase II multicenter study to assess the tolerability and efficacy of the addition of Bevacizumab to standard induction therapy in AML and high risk MDS above 60 years

Lymphoma

SAKK 36/06 | A multicenter phase II trial testing Everolimus (RAD001) for the treatment of patients with relapsed or therapy resistant mantle cell lymphoma

SAKK 37/05 | Ibrutinomab tiuxetan and high-dose melphalan as conditioning regimen before autologous stem cell transplantation for elderly patients with lymphoma in relapse or resistant to chemotherapy. A multicenter phase I trial

SAKK 38/07 | Prospective evaluation of the predictive value of PET in patients with diffuse large B-cell-lymphoma under R-CHOP-14. A multicenter study

CORAL 50-03B | GELARC-CORAL Collaborative Trial in Relapse/Refractory Aggressive Lymphoma Randomized study of ICE plus rituximab (R-ICE) versus DHAP plus rituximab (R-DHAP) in previously treated patients with CD 20 positive diffuse large B-cell lymphoma, eligible for transplantation followed by randomized maintenance treatment with Rituximab

EBMT MMVAR | A Randomized controlled study of Velcade (Bortezomib) plus Thalidomide plus Dexamethasone compared to Thalidomide plus Dexamethasone for the treatment of myeloma patients progressing or relapsing after autologous transplantation

HD13 | Morbus Hodgkin in adults, limited stages

HD14 | Morbus Hodgkin in adults, intermediate stages

HD15 | Morbus Hodgkin in adults, advanced stages

IFM 2005-02 | Relevance of maintenance therapy using Lenalidomide (REVLIMID®) after autologous stem cell transplantation in myeloma patients under the age of 65 (Open, randomised, multi-centric trial versus placebo)

Gastrointestinal Cancer

SAKK 40/04 | Clinical function after total mesorectal excision and rectal replacement. A prospective randomized trial comparing side-to-end anastomosis, colon-J-pouch and straight coloanal anastomosis

SAKK 41/06 | Bevacizumab maintenance versus no maintenance after stop of first-line chemotherapy in patients with metastatic colorectal cancer. A randomized multicenter phase III non-inferiority trial

SAKK 56/07 | Dasatinib first-line treatment in gastrointestinal stromal tumors. A multicenter phase II trial

SAKK 75/06 | Cetuximab in combination with radiation therapy and chemotherapy prior to surgery in patients with resectable, locally advanced esophageal carcinoma. A multicenter phase Ib-II trial

SAKK 77/06 | Continuous sunitinib treatment in patients with unresectable hepatocellular carcinoma. A multicenter phase II trial

SAKK 77/07 | External beam radiotherapy for unresectable hepatocellular carcinoma. A multicenter phase I/II trial

Melanoma

SAKK 50/07 | Temozolomide combined with bevacizumab in metastatic melanoma. A multicenter phase II trial

Sarcoma

EuroEwing 99 | Studie zur Behandlung des Tumors der Ewing-Gruppe

Supportive Care

SAKK 95/06 | A multicenter randomized controlled phase III study of longitudinal electronic monitoring of symptoms and syndromes associated with advanced cancer in patients receiving anticancer treatment in palliative intention

Central Nervous System Cancer

SAKK 70/03 | Whole brain radiotherapy in combination with gefitinib (Iressa) or temozolomide (Temodal) for brain metastases from non-small cell lung cancer (NSCLC). A randomized phase II trial

New Drugs

S065APOX01 | Phase I dose finding and pharmacokinetic study of intravenous APO010, a recombinant form of human Fas ligand, in patients with solid tumors

S065ST1901 | Phase I dose finding and pharmacokinetic study of the intravenous camptothecin ST1968 in patients with solid tumors

SKSD00701 | Dose-finding study of satraplatin in combination with oral vinorelbine in patients with advanced solid tumors. A SAKK-SENDO phase Ib study

SKSD00702 | A phase IB study of the histone deacetylase inhibitor Panobinostat (LBH589) given orally in combination with Carboplatin and Paclitaxel in patients with advanced solid tumors. A SAKK-SENDO phase Ib study

Trials Activated in 2008**Breast Cancer**

SAKK 21/07 | Swiss Observational Bone Study: a substudy of BIG 1-98

Leukemia

SAKK 30/07 | 5-Azacytidine to treat acute myeloid leukemia in elderly or frail patients not suitable for extensive chemotherapy. A multicenter phase II trial

Gastrointestinal Cancer

SAKK 56/07 | Dasatinib first-line treatment in gastrointestinal stromal tumors. A multicenter phase II trial

SAKK 77/07 | External beam radiotherapy for unresectable hepatocellular carcinoma. A multicenter phase I/II trial

Urogenital Cancer

SAKK 08/07 | Docetaxel and cetuximab in patients with docetaxel-resistant hormone-refractory prostate cancer (HRPC). A multicenter phase II trial

Trials closed in 2008**Breast Cancer**

SAKK 24/06 | Bevacizumab and pegylated liposomal doxorubicin as first-line therapy for locally recurrent or metastatic breast cancer. A multicenter, single-arm phase II trial

Closed for accrual on 05.09.2008

Gastrointestinal Cancer

SAKK 75/06 | Cetuximab in combination with radiation therapy and chemotherapy prior to surgery in patients with resectable, locally advanced esophageal carcinoma; a multicenter phase Ib-II trial

Closed for accrual on 28.11.2008

SAKK 76/02 | Docetaxel and cisplatin chemotherapy followed by radiochemotherapy in patients with inoperable, locally advanced esophageal cancer

Closed for accrual on 21.02.2008

SAKK 77/06 | Continuous sunitinib treatment in patients with unresectable hepatocellular carcinoma. A multicenter phase II trial

Closed for accrual on 07.08.2008

Leukemias

SAKK 30/00/HOVON 42 | Randomized induction and post induction therapy in adult patients with acute myelocytic leukemia or refractory anemia with excess of blasts (RAEB, RAEB-t) with IPSS score ≥ 1.5

Closed for accrual on 26.08.2008

Lymphomas

CORAL 50-03B | Randomized study of ICE plus rituximab versus DHAP plus rituximab in previously treated patients with CD20 positive diffuse large B-cell lymphoma, eligible for transplantation followed by randomized maintenance treatment with rituximab

Closed for accrual on 01.07.2008

HD15 | Morbus Hodgkin in adults, advanced stages

Closed for accrual on 18.04.2008

IFM 2005-02 | Relevance of maintenance therapy using Lenalidomide (REVLIMID®) after autologous stem cell transplantation in myeloma patients under the age of 65 (Open, randomised, multi-centric trial versus placebo)

Closed for accrual on 25.08.2008

Project Group Breast Cancer



Presidents:

- 1 Prof. Dr Stefan Aebi, Department of Medical Oncology, Inselspital, University Hospital Bern (until end of 2008)
- 2 PD Dr Georges Vlastos, Department of Gynecology, Breast Unit, University Hospital Geneva (HUG)
- 3 Prof. Dr Christoph Rochlitz, Department of Medical Oncology, University Hospital Basel (starting January 2009)

Objectives

The Breast Cancer Project Group (BCPG) aims to facilitate and conduct clinical and translational research in breast cancer and to collaborate with international research groups (i. e. IBCSG, BIG, EORTC). In the trials SAKK 22/99, 23/03, 24/06, 21/07, and IBCSG 22, 23, 24, 25, 27, 35, 36 and IBIS II, these objectives have been reached. In addition, the BCPG keeps its members updated on clinical trials of IBCSG and BIG, and has reached a high visibility of members of the project group in the breast cancer community, it also cultivates excellent international relations.

The reintegration of the gynecologists, an important objective for the BCPG, is still ongoing. As of 2008, a gynecologist is co-president of the BCPG. The «Arbeitsgemeinschaft für Onkologie», AGO, has agreed not to develop projects in breast cancer and to focus on gynecologic malignancies. Several gynecologists are members of the BCPG.

With the trial SAKK 92/08 the BCPG develops a project in supportive care to prevent PPE (palmar-plantar erythrodysesthesia) in patients indicated for treatment with pegylated liposomal doxorubicin.

Activities

Trial Activated in 2008

SAKK 21/07 | *Swiss Observational Bone Study: a substudy of BIG 1-98*

The primary objective was to construct a statistical model that describes the evolution of bone mineral density in patients treated in each arm of the BIG 1-98 trial.

This model could generate data about BMD modifications over time and potentially be applicable to all patients, rather than a selected substudy population. This will increase the knowledge about the patients who are most «bone-fragile».

The study was activated on June 16, 2008. An abstract has been submitted to ASCO 2009.

Closed Trials

SAKK 24/06 | *Bevacizumab (Avastin®) and pegylated liposomal doxorubicin (Caelyx®) as first-line therapy for locally recurrent or metastatic breast cancer. A multi-center, open-label phase II trial*

The main objective of this trial was to evaluate the safety and tolerability of pegylated liposomal doxorubicin in combination with bevacizumab in patients with inoperable locally recurrent or metastatic breast cancer.

After all patients (43) registered had reached the required observation time of 6 months, the evaluation of the primary endpoint had failed to demonstrate sufficient tolerability of the combined treatment. Therefore, and according to the requirements of the trial protocol, the trial was definitively and permanently closed as of September 5th 2008.

The trial is not yet published, the preparation of the manuscript is planned in 2009.

Strategic Elements for the Next Two Years

In the next two years the group will focus its activities on the following strategic elements:

- To facilitate and conduct clinical and translational research in breast cancer
- To focus on metastatic breast cancer
- To develop non-drug trials
- To collaborate with international research groups
- To define membership in terms of collaboration and responsibilities
- To create team spirit (at least in a core group of active members)
- To provide opportunities for the professional development of young researchers
- To consider the integration of a patient advocate

Portfolio Plan

The portfolio plan for the next years contains the following trials:

SAKK 92/08 PPE trial | *Local antiperspirans for prevention of palmar-plantar erythrodysesthesia (PPE) in patients treated with pegylated liposomal doxorubicin: A randomized, multicenter, double blinded, phase III trial*

The aim of this trial is to evaluate the effects of F511 cream on the occurrence of palmar-plantar erythrodysesthesia (PPE) in patients with breast cancer treated with pegylated liposomal doxorubicin.

The trial was accepted by the SAKK Board on October 21, 2008, and is being developed.

SAKK 21/08 | *Fulvestrant with or without AZD6244, a mitogen-activated protein kinase kinase (MEK) 1/2 inhibitor, in advanced stage breast cancer progressing after first-line aromatase inhibitor: a randomized phase II trial*

The primary objective of the trial is to assess the activity of the combination fulvestrant + AZD6244 in patients progressing after first-line AI.

Project Group Gastrointestinal Cancer



President:

Prof. Dr Markus M. Borner, Clinical Research Unit of the Oncology Department, Inselspital, University Hospital Bern and Oncology Unit, Hospital Biel

Objectives

The main objectives of the project group reached since 2007 have been to cover as many clinical situations in Gastrointestinal cancer as possible with interesting protocols, to avoid competition with registration studies of the industry, as well as to promote the international collaboration.

Activities

Trials Activated in 2008

SAKK 56/07 | *Dasatinib first-line treatment in gastrointestinal stromal tumors. A multicenter phase II trial*

The primary objective is to assess the efficacy of dasatinib treatment by fusion PET/CT-scan.

Secondary objectives are to assess the efficacy and safety of dasatinib in GIST, to correlate efficacy of dasatinib with KIT and PDGRF mutation status, the efficacy and safety with dasatinib drug exposure, and to assess the efficacy of second-line treatment with another TK-inhibitor.

The trial was activated on January 18, 2008.

SAKK 77/07 and SASL 26 | *External beam radiotherapy for unresectable hepatocellular carcinoma. A multicenter phase I/II trial*

HCC is the fifth most common cancer worldwide with increasing incidence rates in Europe and high mortality rates. The standard care for non-resectable liver cancer is TACE. Alternative methods are radiofrequency ablation (RFA), nuclear therapy with intra-arterial radioactive beads and ionizing radiation. Ionizing radiation is a highly effective single agent against HCC. Radiotherapy has been underused in the past due to technical inadequacy. More recent technical developments such as image-guided radiotherapy, CT-based RT planning, and 3D-

conformal dose delivery allow minimizing the dose to non-target tissue such as non-affected liver tissue, kidneys and the gut. Thus, less collateral dose delivery to healthy neighboring tissue allows dose escalation to the diseased segments. The present trial is the first to investigate and define the potential role of high-dose external beam RT for non-resectable HCC in Switzerland and a few centers abroad, seeking to establish RT as a potential experimental arm in a subsequent phase III trial.

The trial was activated on November 7, 2008.

Closed Trials

SAKK 75/06 | *Cetuximab in combination with radiation therapy and chemotherapy prior to surgery in patients with resectable, locally advanced esophageal carcinoma. A multicenter phase Ib-II trial*

The trial was closed on November 28, 2008. There was no unexpected toxicity observed. In 12 centers 28 patients were enrolled.

SAKK 76/02 | *Docetaxel and cisplatin chemotherapy followed by radiochemotherapy in patients with inoperable, locally advanced esophageal cancer. A multicenter phase II trial*

At the interim analysis 21 patients were evaluated, of which four had local control, instead of the eight required for the trial continuation. Some patients were not evaluated according to protocol, since distant progression was documented and endoscopy and biopsies were not considered clinically relevant. The trial thus is formally negative. The therapy schedule was however well tolerated, with only 3/21 patients having e. g. grade III dysphagia under RCT. The outcome in this rather poor patient group was acceptable, particularly since some patients with residual tumor cells on biopsy had clinically meaningful long-term control. The results were presented in a poster at the ESMO 2008 meeting.

In seven centers 21 patients were enrolled. The trial was closed on February 21, 2008.

SAKK 77/06 | *Continuous sunitinib treatment in patients with unresectable hepatocellular carcinoma. A multicenter phase II trial*

The interim analysis included the first 24 of the total 45 patients required for this trial as stated in the trial protocol. No protocol violations were committed regarding trial inclusion/exclusion for the first 24 patients. The primary endpoint was progression-free survival

12 weeks after trial enrollment. The observed number of patients who were alive and progression-free at 12 weeks satisfied the criterion for trial continuation. No cause for concern was raised with regard to (serious) adverse events.

Due to a fast accrual, this trial was closed earlier than expected on August 7, 2008. In nine Swiss centers 31 patients were enrolled, in one center in Budapest 14 patients.

Strategic Elements for the Next Two Years

In interdisciplinary trials the focus will be brought on

- the treatment sequence in synchronous CRC
- Intraperitoneal chemotherapy
- Radiation vs other options in HCC
- Favor clear-cut translational questions over therapeutic fishing expeditions

Each trial should

- have one major translational endpoint
- consider aspects of quality, infrastructure and available resources in Switzerland

Portfolio Plan

The following trials are in the process of discussion or activation:

SAKK 41/07 | *Neoadjuvant treatment with radiotherapy and capecitabine with or without panitumumab in patients with advanced, K-ras unmutated rectal cancer. A randomized multicenter phase II trial*

The objectives of this trial are to assess the efficacy and safety of the neoadjuvant regimen of capecitabine and external beam radiotherapy (RT) in combination with panitumumab in patients with advanced K-ras unmutated rectal cancer.

SAKK 41/08 | *Neoadjuvant radiotherapy combined with capecitabine and sorafenib in patients with advanced, K-ras mutated rectal cancer. A multicenter phase I/IIa trial*

The objective of the phase I part is to determine the recommended dose of the neoadjuvant regimen of capecitabine, sorafenib and external beam radiotherapy (RT) in patients with advanced K-ras mutated rectal cancer.

The objectives of the phase IIa part are to assess the efficacy and safety of the neoadjuvant regimen of capecitabine, sorafenib and external beam RT in patients with advanced K-ras mutated rectal cancer.

SAKK 77/08 | *Sorafenib alone or in combination with everolimus in patients with unresectable hepatocellular carcinoma. A randomized multicenter phase II trial*

The objective of the trial is to investigate if the combination of sorafenib plus everolimus can stop tumor progression, with a sorafenib monotherapy group used to control selection bias.

SAKK 77/09 | *A phase I open label/Phase II randomized, double-blind, multicenter trial investigating the combination of Everolimus and TransArterial ChemoEmbolisation (TACE) with doxorubicin in patients with hepatocellular carcinoma eligible for TACE*

The objective of the phase I part is to determine the maximum tolerated (MTD) dose and dose limiting toxicities of the administration of daily everolimus in patients with HCC treated with TACE.

The objective of the phase II part is to determine the efficacy and tolerability of the administration of daily everolimus in patients with HCC treated with TACE as compared to TACE alone.

NCIC BI | *A phase III study of gemcitabine plus capecitabine (GEMCAP) versus gemcitabine alone in advanced biliary cancer*

The primary objective of this study is to compare the overall survival in patients with histologically or cytologically proven adenocarcinoma of the biliary tree that are either unresectable or metastatic and randomized in a 1:1 fashion to receive either gemcitabine and capecitabine (GEMCAP) or gemcitabine alone (GEM) as first-line chemotherapy. The progression-free survival, response rate, duration of response, rate of stable disease, rate of disease control, toxicity and quality of life between the two treatment arms will also be compared. The relationship between the expression of baseline tumor molecular marker (at diagnosis) with outcomes and response to treatment will also be explored.

SAKK 75/08 | *Multimodal therapy with and without cetuximab in patients with locally advanced esophageal carcinoma. Unblinded, prospectively randomized phase III trial*

The primary objective of the trial is to determine the efficacy of neoadjuvant chemoradiotherapy (CRT) combined with immunotherapy followed by adjuvant immunotherapy compared with the same schedule without immunotherapy (neoadjuvant and adjuvant).

Secondary objectives of the trial are to compare the toxicity of the two therapy arms and to determine duration of response and patterns of failure.

EORTC 40071 | An EORTC trial on the role of lapatinib in advanced gastric cancer is in development in the Gastrointestinal Group under the lead of PD Dr Arnaud Roth.

Collaboration with/Participation in Other Groups

The Project Group Gastrointestinal Cancer maintains the collaboration with other SAKK groups. The established collaboration with foreign Groups/Centers e.g. the German esophageal cancer group, EORTC, NCIC and Italian centers (EIO) is continued.

Project Group Leukemia



President:

PD Dr Yves Chalandon, Hematology Service,
University Hospital Geneva (HUG)

Objectives

We offer clinical studies covering the main topics in acute and chronic leukemia, however not myelodysplasia (MDS) low risk and Myeloproliferative Disorders (MPD). The project group collaborates with international study groups in developing and performing phase III trials. But still, more participation of Swiss members in international cooperative groups is desirable. Phase I–II trials testing new compounds and combinations are being developed. The project group also participates in international working groups. We have established a platform for younger clinical researchers, and some younger investigators are now involved in SAKK trials. Furthermore, the project group plans the foundation of a Swiss registry for acute leukemia, and will check to take over the lead in Phase III trials. The objective to have active membership working in the field of acute and chronic leukemia has been partially achieved as still too few members are active (around 10–15). With the smaller centers participating in SAKK we are hoping to have more success in including patients particularly in chronic leukemia trials (partially achieved) in all Swiss centers.

Activities

Trials Activated in 2008

Phase III trials:

APL 2006 (Acute Promyelocytic Leukemia) | *Randomized phase III trial assessing the role of arsenic trioxide and/or ATRA during consolidation course in newly diagnosed acute promyelocytic leukemia (APL)*

We hope that the event-free survival at two years will further increase at the investigational arms with reduced toxicity and without increasing the relapse rate by comparison with a classical anthracycline-AraC consolidation regimen. The total accrual target is 895 patients. The trial was activated on April 8, 2008.

Phase II trials:

AML (Acute Myeloid Leukemia)

SAKK 30/07 | *5-Azacytidine to treat acute myeloid leukemia in elderly or frail patients not suitable for intensive chemotherapy. A multicenter phase II trial*

The main objective of the trial is to evaluate the efficacy of 5-azacytidine in a population of patients with AML not suitable for induction type chemotherapy because of age or relevant comorbidities. Secondary objectives are to evaluate survival and adverse events. The trial was activated on September 23, 2008.

Translational research

SAKK 63/03 | *Blood and bone marrow banking in SAKK leukemia trials*

The main objective of the trial is to preserve material for later use in biological studies which will be submitted to SAKK in the future. The members of the SAKK Leukemia Project Group have decided to bank the material centrally in Aarau. It will be collected at the time of inclusion of a patient into one of the ongoing trials. The project is supervised by a banking committee. The database documenting the collection and central storage of material resides at the SAKK Coordinating Center. A remote data entry facility has been developed to this effect. It will also provide to the researchers an overview of the banked material, thus enabling them to check if enough material is available for a potential future study. 61 samples have been collected.

Closed Trials

Acute Myeloid Leukemia, AML/high risk MDS

SAKK 30/00 | *Randomized induction concerning the value of G-CSF priming in adult patients (≤ 60 yrs of age) with acute myelocytic leukemia (AML) or refractory anemia with excess of blasts (RAEB, RAEB-t) with IPSS score ≥ 1.5*

The trial was testing the value of priming leukemic cells with G-CSF during induction and post-induction chemotherapy in patients ≤ 60 years. Remarkable for such a long-term trial, the accrual continued to be as planned.

The trial was closed on August 26, 2008, after having recruited a total of 154 AML patients among the Swiss centers. No major protocol violations have been reported.

Strategic Elements for the Next Two Years

- to develop phase II trials for patients with acute leukemia unfit for intensive chemotherapy or for elderly patients with new drugs targeted therapy (in combination with low-dose sequential chemotherapy) or vaccines
- to stimulate translational research projects (prognostic MRD (Minimal Residual Disease) as well as study of leukemic stem cells, leukemogenesis, genomic and proteomic) as this was poorly done for the last two years. We need to have more collaboration with research laboratories.
- to improve the input of SAKK in the collaboration with international study groups as far as clinical phase III trials are concerned
- to create a Swiss registry for acute leukemia

Portfolio Plan

Trials

Phase III:

CLL 10 protocol of the German CLL Study Group (GCLLSG) |
Phase III trial of combined immunochemotherapy with Fludarabine, Cyclophosphamide and Rituximab (FCR) versus Bendamustine and Rituximab (BR) in patients with previously untreated chronic lymphocytic leukemia
 The trial will be activated in 2009.

EBMT RIC-MUD AML | *A Randomized Phase III study comparing conventional chemotherapy to low-dose total body irradiation-based conditioning and hematopoietic cell transplantation from related and unrelated donors as consolidation therapy for older patients with AML in first Complete Remission*
 The trial will be activated in 2009.

Phase II:

SAKK 32/09 | *Phase II trial of sequential administration of Rituximab and Cladribine in the treatment of patients with newly diagnosed, relapsed and refractory hairy cell leukemia and variant forms.*
 The SAKK Board will decide about the conduct of the trial in the final assessment in 2009.

Phase I:

SAKK 65/08 | *In collaboration with the Phase I project group and the lymphoma project group Nelfinavir and Bortezomib in patients with hematological malignancies who are not eligible for intensive cytotoxic therapy*
 The trial proposal is being discussed and has to be approved by the SAKK Board.

Follow-up Trials

HOVON 92/SAKK 30/08 (follow-up HOVON 42/SAKK 30/00) |
A phase II-III randomized study to assess the added value of Laromustine in combination with standard remission-induction chemotherapy in patients aged 18-65 years with previously untreated acute myeloid leukemia (AML) or myelodysplasia (MDS) (RAEB with IPSS \geq 1.5)
 The trial will be activated in 2009.

CML V, Chronic Myeloid Leukemia with the German Study Group, which should follow the CML IV protocol
 The trial is under discussion in the project group.
 A follow-up trial of the SAKK 30/07 AML trial for frail elderly AML patients.

Primary objective: to test the feasibility and efficacy of combination therapy of 5-Azacytidine with another new drug (AML targeted therapy)


The trial is under discussion in the project group.

Collaboration with/Participation in Other Groups

The Leukemia Project Group collaborates with the Lymphoma Project Group within the SAKK and with the following other groups:

- Project Group New Anticancer Drugs
- Laboratory group (molecular diagnostic), Swiss Molecular Hematology/Oncology Working Group
- The Dutch HOVON group in AML
- The collaborative group GRAALL (Group for Research in Adult Acute Lymphoblastic Leukemia) including the French groups GOELAMS-LALA, Belgium in ALL
- The German CLL Study Group (GCLLSG) in CLL
- The German CML Study Group (GCMLSG) in CML
- The European APL group
- The European Leukemia Network
- The European Group for Blood and Marrow Transplantation (EBMT)

Project Group Lung Cancer



Presidents:

- 1 PD Dr Miklos Pless, Department of Medical Oncology and Tumor Center, Kantonsspital Winterthur
- 2 Prof. Dr Walter Weder, Division of Thoracic Surgery, University Hospital Zurich

Objectives of the Lung Cancer Project Group

- The Lung Cancer Project Group offers interesting studies to Swiss cancer centers to treat as many Non-Small Cell Lung Cancer (NSCLC) patients as possible in trials (stage IV).
- It establishes a network of Swiss lung cancer centers with multidisciplinary thoracic capacity (stage IIIB/IIIA), and a solid basis for translational research (tissue banking).
- The group has become an attractive partner for pharmaceutical companies with interesting compounds, and helps to advance the career of young oncologists.

Activities

Open Trials in 2008

There were no new trials activated in lung cancer in 2008, but we made important progress in SAKK 16/00, SAKK 17/04, and SAKK 19/05.

Strategic Elements for the Next Two Years

- to establish follow-up studies for stage IV (SAKK 19/09), stage IIIB (SAKK 16/08) and also open a new Small-Cell Lung Cancer trial (SAKK 15/08)
- to establish a cooperation with other cooperative groups, e.g. the Belgian group in Leuven
- to support the ETOP platform
- to publish translational research data on our phase II trials

Portfolio Plan

SAKK 15/08 SCLC, Small-Cell Lung Cancer | *Carboplatin and Paclitaxel plus ASA404 as first-line chemotherapy for extensive-stage small-cell lung cancer (SCLC). A phase II trial*

The main objective is the efficacy and feasibility of this combination in SCLC.

The trial is resubmitted to the SAKK Board.

SAKK 16/08, NSCLC | *Preoperative chemo-radiotherapy combined with concomitant Cetuximab in non-small cell lung cancer (NSCLC) patients with IIIB disease. A multicenter phase II trial*

The objective of this trial is to evaluate activity and safety of sequential neoadjuvant chemo-radiotherapy with concomitant targeted therapy of cetuximab in operable stage IIIB NSCLC patients.

The protocol is being developed and will be submitted to the SAKK Board.

SAKK 19/09, NSCLC | *Pemetrexed, cisplatin and bevacizumab, or erlotinib and bevacizumab for metastatic adenocarcinoma of the lung according to EGFR mutation status: a multicenter phase II study including biopsy at progression (BIOPRO trial)*

The trial proposal has to be approved by the SAKK Board.

Collaborations with Working Groups, Sections and Other Groups:

- Leuven (Belgium)
- Novi Sad (Serbia)

Project Group Lymphoma



President:

PD Dr Nicolas Ketterer, Centre Pluridisciplinaire d'Oncologie, University Hospital Lausanne (CHUV)

Objectives

The Lymphoma Project Group's main objectives are to bring together onco-hematologists and other specialists involved and interested in the management of lymphoma/myeloma patients, to ameliorate the management and the treatment of patients with lymphoma, by developing and leading some original clinical trials. The project group stimulates and promotes translational research for a better understanding of lymphoid malignancies, with the aim to improve the treatment of the patients in the following projects:

- Minimal Residual Disease, MRD and other ancillary studies in follicular lymphoma (FL) trials
- MRD + Ig-VH status in mantle cell lymphoma trials
- Participation to the pathological subprojects in the CORAL trial

Activities

Open Trials in 2008

There were no new trials activated in lymphoma in 2008, but we made important progress in SAKK 36/06, SAKK 37/05, SAKK 38/07 and HD13, HD14, and EBMT MMVAR.

Closed Trials

IFM 2005-02 | Intergroupe Francophone du Myélome (maintenance after Autologous Stem Cell Transplantation, ASCT)

Relevance of maintenance therapy using Lenalidomide (REVLIMID®) after autologous stem cell transplantation in myeloma patients under the age of 65 (Open, randomized, multi-centric trial versus placebo)

International, open, phase III trial (Revlimid® versus placebo) comparing the direct clinical benefit.

The trial has been closed for accrual in August 2008 after the planned sample size had been reached. Twenty-five patients have been enrolled in eight centers in Switzerland.

HD15 | *Morbus Hodgkin in adults, advanced stages*

Reducing toxicity by de-escalation, while maintaining efficacy

This trial has been activated in 13 centers in Switzerland.

CORAL study (ASCT for DLBCL in 1st relapse) | GELARC-CORAL

Collaborative Trial in Relapse/Refractory Aggressive Lymphoma. *Randomized study of ICE plus rituximab (R-ICE) versus DHAP plus rituximab (R-DHAP) in previously treated patients with CD20 positive diffuse large B-cell lymphoma, eligible for transplantation followed by randomized maintenance treatment with Rituximab*

The trial has been closed last summer. Twenty-four patients have been enrolled in 10 centers in Switzerland. The second randomization of the last patient has been foreseen in December 2008. The end of maintenance for the last patient should be reached in December 2009. An analysis for R-ICE versus R-DHAP on the first 400 patients is in preparation.

Strategic Elements for the Next Two Years

The project group adjusts its activities to comply with the objectives requested by the Scientific Advisory Board in 2007. The group concentrates on key trials and therefore, will build on pre-existing collaborations and launch a key clinical trial under SAKK's guidance. Furthermore it plans to participate as SAKK member in major European trials.

Portfolio Plan

Diffuse Large B-Cell-Lymphoma, DLBCL

SAKK 38/08 (relapsed elderly) | *Rituximab, bendamustine and lenalidomide in elderly and/or frail patients with relapsed or refractory aggressive non-Hodgkin's Lymphoma. A phase III trial*

The objective of the phase I part is to determine the MTD (The highest dose of a drug treatment that does not cause unacceptable side effects) of the combination of rituximab, bendamustine and lenalidomide in patients with relapsed refractory aggressive Non-Hodgkin Lymphoma, NHL, not eligible for high-dose therapy with ASCT.

The objective of the phase II part is to determine the efficacy and safety of the combination of rituximab, bendamustine and lenalidomide in patients with refractory or relapsed aggressive NHL not eligible for high-dose therapy with ASCT.

The trial has been accepted by the SAKK Board and is being developed.

Follicular Lymphoma, FL

SAKK 35/09 | *GA 101 versus Rituximab for untreated follicular lymphoma. An intergroup randomized phase II trial*

The primary objective of this trial is to investigate if GA 101 is more effective than rituximab. The trial will be developed in collaboration with the Nordic Lymphoma Group.

Multiple Myeloma, MM

IFM 2009 trial | *Randomized phase III study comparing conventional dose treatment using a combination of Bortezomib, Lenalidomide and Dexamethasone to high-dose treatment with peripheral stem cell transplant in the initial management of myeloma in patients under 65 years of age*

The aim of the protocol is to determine if, in the era of new drugs, high-dose treatment is still necessary in the initial management of myeloma in young patients. The primary objective is to compare event-free survival in the two arms. The trial will be activated in France at the end of 2009.

SAKK 65/08 | *Synergistic targeting of the ER stress response with Nelfinavir and Bortezomib: a phase I dose escalation trial in advanced hematologic malignancies*

The aim of the trial is to establish tolerability and toxicity of the induction of UPR (unfolded protein response) activity and/or the inhibition of akt-phosphorylation by nelfinavir in combination with bortezomib in patients with advanced hematopoietic malignancies, and to establish a recommended dose for a phase II trial.

This trial will be developed and conducted in collaboration with the Project Group Leukemia and with the Phase I Project Group. Patients will be accrued in selected centers. The trial has to be approved by the SAKK Board in 2009.

HD18 | *Treatment optimization in the first-line treatment of advanced stage Hodgkin lymphoma; treatment stratification by means of FDG-PET*

The aim of the trial is to individualize treatment for each patient by adapting it to early response and thus to continue intensive treatment only with those patients who show an inadequate treatment response.

The trial has been activated in Germany on May 13, 2008 and is open for accrual in Germany. The trial will be activated 2009 in Switzerland.

Collaborations

The Project Group seeks to have an active role in selected large international collaborative projects, allowing a reinforcement of our collaboration with other large cooperative international groups as:

- German Hodgkin Study Group (HD trials)
- Intergroupe Francophone du Myélome (IFM trials)
- GELA, (CORAL study)
- European Mantle Cell Lymphoma Network

Project Group New Anticancer Drugs/ Phase I Trials



President:

PD Dr Cristiana Sessa, Oncology Institute of Southern Switzerland (IOSI) Bellinzona

Objectives

The primary aim of the project group is to increase the active participation in Phase I trials and to get new drugs to be tested by SAKK in Phase II trials; the group also aims to increase experience and set up a central coordination for early drug development.

SAKK and SENDO have established a collaboration in order to increase and improve the involvement of selected SAKK centers in early clinical trials, and to provide SAKK with a constant flow of new drugs for Phase II trials.

Activities

Trial Activated in 2008

SKSD00702 | *A phase Ib study of the histone deacetylase inhibitor Panobinostat (LBH589) given orally in combination with Carboplatin and Paclitaxel in patients with advanced solid tumors. A SAKK-SENDO phase Ib study*

The primary objective of the trial is to determine the Maximum Tolerated Dose (MTD) of panobinostat (LBH589) when administered in combination with carboplatin and paclitaxel in patients with advanced solid malignancies and to identify the Recommended Dose (RD) for a subsequent phase II trial.

The trial was activated on May 20, 2008.

Participating centers: Bellinzona (IOSI), Lausanne (CHUV), Basel (USB)

Portfolio Plan

SAKK 65/08 | *Synergistic targeting of the ER stress response with Nelfinavir and Bortezomib: a phase I dose escalation trial in advanced hematologic malignancies*

This trial is developed in collaboration with the Project Group Leukemia and the Project Group Lymphoma. Patients will be accrued in selected centers. The trial is still in discussion and has to be approved by the SAKK Board in 2009.

Collaboration with/Participation in Other Groups

- Project Group Breast Cancer
- Project Group Leukemia
- Project Group Lymphoma
- SENDO Southern Europe New Drugs Organization

Project Group Urogenital Tumors



Presidents:

- 1 PD Dr Silke Gillissen, Department of Medical Oncology, Kantonsspital St. Gallen
- 2 Prof. Dr George Thalmann, Department of Urology, Inselspital, University Hospital Bern

Objectives

Since the group was promoted from a working group to a project group, it has reached a better attendance from urologists, pathologists, and oncologists, and recently also from radio-oncologists.

A new culture of collaboration with the urologists was developed and is ongoing, especially in the STAMPEDE trial and a planned bladder cancer trial.

Ongoing, good translational research opportunities with Prof. Holger Moch, University Hospital Zurich, and Prof. Lukas Bubendorf, University Basel, and Prof. Wilhelm Krek, ETH Zurich, are realized in the development of projects in prostate cancer.

Activities

Trial Activated in 2008

SAKK 08/07 | *Docetaxel and cetuximab in patients with docetaxel-resistant hormone-refractory prostate cancer (HRPC). A multicenter phase II trial*

The main objectives of this trial are to assess the efficacy and safety of the chemo-immunotherapy treatment with docetaxel and cetuximab in docetaxel-resistant patients with HRPC (second-line therapy in prostate cancer).

The trial was activated on June 17, 2008.

Closed Trials

None

Strategic Elements for the Next Two Years

The Project Group Urogenital Cancer aims

- to pursue trials in prostate cancer as main topic
- to involve more centers
- to develop second-line therapies
- to develop therapies in indolent PSA (Prostate-Specific Antigen) rising conditions (metastatic or not) despite androgen ablation
- to conduct trials in palliative care
- to strengthen the collaboration between oncologists, radio-oncologists and urologists

Portfolio Plan

Trials:

SAKK 08/08 | *RAD001 in Hormone Refractory Prostate Cancer as first-line treatment*

The main objectives of this trial are to assess the efficacy and safety of everolimus as first-line therapy in non-rapidly progressive CRPC (Castration-Resistant Prostate Cancer).

SAKK 08/08 is in a protocol finalizing state.

STAMPEDE | *Systemic Therapy in Advancing or Metastatic Prostate Cancer: Evaluation of Drug Efficacy. A 5-stage multi-arm randomised controlled trial*

The trial will be activated in Swiss centers in 2009.

Follow-up Trials:

- A second-line trial in prostate cancer is planned.
- If SAKK 08/07 is successful, a Phase III trial in first-line HRPC (Docetaxel/Cetuximab) will be planned.

Collaboration with/Participation in Other Groups

- Collaboration with other SAKK groups such as PG New Anticancer Drugs
- Collaboration with foreign Groups/Centers like MRC (Medical Research Council)
- Collaboration with other Cooperative Groups such as EORTC (European Organisation for Research and Treatment of Cancer)

Section Clinical Research Coordinators (CRC)



Presidents:

- 1 Christine Biaggi Rudolf, SAKK Coordinating Center Bern
- 2 Julia Rengier-Styles, Centre Pluridisciplinaire d'Oncologie, University Hospital Lausanne (CHUV)

Short introduction

From the very beginning the skills required for a position as a Clinical Research Coordinator (CRC) could only be learned on the job. Most people applying for such a position had some background in the medical field since they came from nursing or had worked as a receptionist in a doctor's office before. Nowadays, this position is more and more filled by new PhD graduates who want to acquire their first work experience outside academia after successfully finishing their thesis. Either way, very few have previous experience as a CRC. That is why many years ago the Section of Clinical Research Coordinators was founded, which in the early days was called the Data Managers Section. The goal was to form an interactive group with participants from all the various SAKK centers in Switzerland and from SAKK affiliate centers in the neighboring EU countries that would meet on a regular basis to exchange ideas and experiences on how to run and coordinate data management in clinical trials.

Activities 2008

In January we had our two-day annual meeting for CRCs at the SAKK Coordinating Center in Bern. It was a very intense program with a lot of interesting presentations in various fields of cancer research as well as some excursions into Good Clinical Practice (GCP). The first morning of those two days was particularly intended for new section members who had only recently started their work as a CRC at a center. Overall 45 certificates were handed out indicating the presence of the attendees.

In November the Section met at the semi-annual meeting in Basel. This meeting focused mainly on new procedures (such as: changes regarding documentation of adverse events reported on case report forms, availability of a new pregnancy reporting form) and updates (e. g. on certain aspects of GCP, announcement of introduction of electronic data capturing for SAKK trials in 2009, activation and closure of SAKK trials throughout 2008) presented by SAKK staff on one hand, on the other hand the Clinical Trial Unit (CTU) Basel introduced its educational program for clinical research professionals which is planned to start in fall 2009.

Outlook

In 2009 the section intends to concentrate on finding ways in which we can improve sharing professional knowledge. There are quite a few medium-sized centers, which employ only one CRC – the CRCs working in such a center need support from well organized CRC personnel.

Further, we hope to strengthen the collaboration and discussions between the SAKK and the CTU Bern in order to launch our own educational program for clinical research professional training tailored especially towards CRCs.

In 2009 we will again organize our two-day annual meeting (January 26/27) as well as a section meeting in Basel on November 26 as part of the SAKK semi-annual meeting.

Section Internal Medicine

President:

Dr Rudolf Morant, Department of Internal Medicine,
Tumorzentrum ZeTuP, St. Gallen

Activities 2008

The Section Internal Medicine was working on a new proposal for a phase III trial concentrating on the effect of low-molecular weight heparin (LMWH) on survival in cancer patients with a high risk for thromboembolism.

SAKK 91/08 | *The effect of low-molecular weight heparin (e. g. Enoxaparin) on survival and thromboembolism in a high-risk subgroup of cancer patients.*

The primary objective of this multicenter phase III trial is to demonstrate that LMWH prophylaxis over six months will result in a significantly reduced death rate (25 % relative risk reduction) in a subcategory of cancer patients with a high risk of venous thromboembolism, or VTE.

Secondary objectives are to evaluate the VTE rate and bleeding complications.

SAKK 91/08 would be the first proof of principle trial with LMWH in a high-risk cancer population with a simple, safe and easily available intervention and a hard single endpoint (mortality).

Principal investigators of SAKK 91/08 are Dr Stefanie Pederiva and Prof. Jürg Beer, both at the Kantonsspital Baden, who presented their trial proposal to the project groups Lung Cancer, Gastrointestinal Cancer and Breast Cancer as well as to the section meeting Internal Medicine during SAKK's last semi-annual meeting in Bern. Currently, the centers have to indicate their interest and prospective accrual for this trial. The trial proposal has been re-submitted to the SAKK Board. So far, no pharmaceutical company is willing to provide the drug for this interesting project.

Section Pathology



President:

Prof. Dr Holger Moch, Institute of Surgical Pathology,
Department Pathology, University Hospital Zurich

Short Introduction

The Section Pathology is active in the following areas:

- The quality and the outcome of many clinical trials frequently require input by pathologists.
- Some trials need a review of the initial pathological diagnosis. The goal of such a review is quality assurance and control. Such reviews have become crucial, particularly with targeted therapy, where the most accurate determination of both the tumor type and the selected target is mandatory.
- Most predictive parameters of clinical responses to novel drugs are performed on tumor tissues. Such predictive tests, e. g. k-RAS testing in colorectal cancer are frequently performed in oncological trials. Quality assurance is mandatory in such predictive tests.
- Translational research consists of exploratory investigations of the fundamental mechanisms, which may yield new evidence on predictive or prognostic factors. It may involve clinical and laboratory investigations on clinical material collected during clinical trials. Translational researchers in pathology may screen patients' tissue samples for markers and correlate them with clinical findings with the aim to identify new therapy modalities and novel diagnostics for tumor diseases.
- For translational research projects, the collection of biomaterial is necessary. Pathologists are involved in prospective and retrospective collection of biomaterial and the establishment of biobanks.

Activities 2008

The Section Pathology is involved in many SAKK trials. The section members also play an important role in the activities of the IBCSG, both on a practical level by contributing patients and on an intellectual and leadership level. Fur-

ther, section members continue to enroll patient material in earlier studies and in new SAKK trials. There are many activities in collaboration with the groups Lung Cancer (SAKK 16/00, SAKK 17/04 etc.), Lymphoma (SAKK 38/07), Melanoma (SAKK 50/07) and others. These activities include the collection of biomaterial, translational research and predictive tests. The completion of patient forms (p-form) requires the engagement of many pathologists.

Outlook

The new strategy of SAKK aims to improve the activities of pathologists in different oncology trials. Pathologists will be involved in the early phases of protocol development. In these phases, laboratory tests can be discussed and implemented in the protocols. From an administrative and logistic point of view, improvement of budgeting, implementation and monitoring of pathology activities in clinical trials are the main objectives for 2009. Costs for biomarker research are only partly paid by the SAKK. Therefore, pathologists should perform more extensive exploratory laboratory-based translational research covered by other, third party funding (e.g. from Oncosuisse or the Swiss National Science Foundation SNF). In the last year, the first SNF grant application has been funded on the basis of material collected in a SAKK trial.

Section Radio-Oncology



Presidents:

- 1 PD Dr Ludwig Plasswilm, Department of Radiation Oncology, Kantonsspital St. Gallen
- 2 Dr Sabine Balmer Majno, Division Radio-Oncology, University Hospital Geneva (HUG)

Activities 2008

In 2008 the members of the SAKK Section Radio-Oncology focused on possibilities to increase the activities and participation of radiation-oncology within SAKK protocols.

During the meeting of the Section Radio-Oncology in November 2007 the idea of a new protocol panel of Swiss Radiation Oncology was introduced (Swiss Trials in Radiation Oncology «SWITRO»). This panel should become a platform of cooperation between the Scientific Association of Swiss Radiation Oncology (SASRO) and the Section Radio-Oncology. A detailed proposal for the generation of such a platform to boost clinical research in radiation-oncology in Switzerland has been discussed within the SASRO executive committee and at the semi-annual meetings in 2008. Finally, the majority of the members of the Radio-Oncology Section agreed to accept SWITRO being the new protocol panel of the Radio-Oncology Section. The start-up meeting of the new panel was held on November 11, 2008, and was attended by D. Aebersold, S. Bodis, L. Plasswilm, D. Weber, K. Zaugg, F. Zimmermann (in the future a representative of the Swiss Society of Radiobiology and Medical Physics will also be invited to join the panel). At that meeting 17 proposals of new protocols were discussed and preliminarily ranked by priority for further discussions and activities within SAKK.

Proposals of new trials

- Intensity-modulated radiation therapy (IMRT) + chemotherapy in anal cancer
- Adjuvant radiation therapy in node positive prostate cancer
- Radiochemotherapy plus EGFR-antibody therapy in unresectable recurrent head and neck cancer
- Postoperative radiochemotherapy plus EGFR-antibody therapy in resectable recurrent head and neck cancer
- FET-PET based planning in primary glioblastoma
- Dose escalation in radiation therapy of biochemically relapsed prostate cancer

In addition and as a complementary effort to «SWITRO» members of our section are also involved in proposals of new protocols which are under discussion within the different SAKK project groups and working groups. Besides these activities members of the Radio-Oncology Section are involved in several ongoing or recently finished protocols.

SAKK – Ongoing trials with strong links to the Radio-Oncology Section

SAKK 70/03 | *Whole brain radiotherapy in combination with gefitinib (Iressa®) or temozolomide (Temodal®) for brain metastases from non-small cell lung cancer (NSCLC). A randomized phase II trial*

SAKK 76/02 | *Docetaxel and cisplatin chemotherapy followed by radiochemotherapy in patients with inoperable, locally advanced esophageal cancer. A multicenter phase II trial*

SAKK 75/06 | *Cetuximab in combination with radiation therapy and chemotherapy prior to surgery in patients with resectable, locally advanced esophageal carcinoma. A multicenter phase Ib-II trial*

SAKK 77/07 and SASL 26 | *External beam radiotherapy for unresectable hepatocellular carcinoma. A multicenter phase I/II trial*

SAKK 16/00 | *Preoperative chemoradiotherapy vs. chemotherapy alone in non-small cell lung cancer (NSCLC) patients with mediastinal lymph node metastases (stage IIIA, N2): A randomized prospective phase III trial*

SAKK 17/04 | *Neoadjuvant chemotherapy and extrapleural pneumonectomy of malignant pleural mesothelioma (MPM) with or without hemithoracic radiotherapy. A randomized multicenter phase II trial*

SAKK Trials in cooperation with the German Hodgkin Lymphoma Study Group

HD13, Limited stages | *Qualitätssicherungsprotokoll zur Toxizitätsreduktion in der Primärtherapie des frühen Morbus Hodgkin*

HD14, Intermediate stages | *Qualitätssicherungsprotokoll zur Effektivitätssteigerung in der Primärtherapie des intermediären Morbus Hodgkin*

Outlook

In 2009 the Radio-Oncology Section will mainly focus on the initiation of new SAKK protocols, based on our current proposals, and to support the ongoing SAKK 77/07 trial. Also the cooperation of the Radio-Oncology Section and SASRO will be advanced. Special thanks go to Daniel Aebbersold, Inselspital Bern, and president of SASRO, who strongly supports this cooperation. Daniel Aebbersold and Ludwig Plasswilm together were personally strongly involved into the set-up and organization of our new joint protocol panel «SWITRO».

Section Surgery



Presidents:

- 1 Prof. Dr Walter Richard Marti, Department of Surgery, University Hospital Basel (until end of January 2009)
- 2 Prof. Dr Paul M. Schneider, Department of Surgery, University Hospital Zurich (starting February 2009)

Short Introduction

The Section Surgery aims to design and conduct clinical and translational research in the field of surgically treated cancers and functions as a platform to promote other multicenter trials in the surgical community.

Activities 2008

The Section Surgery has held its annual meeting on November 20, 2008, in Basel. The main focus of activities in 2008 was to continuously work on the SAKK 40/04 trial. The primary objective of this trial is to compare a 5 cm colon-J-pouch, side-to-side anastomosis and straight colo-anal anastomosis as reconstruction techniques after rectal cancer surgery with respect to defecation quality (evacuation problems, fragmentation of stools), stool frequency, mortality and morbidity. As secondary objectives, the three different reconstruction techniques are compared in respect of short and mid-term quality of life and long-term clinical outcome. As the patient accrual is favorable (~50/year) but not as high as expected (94/year), we still encourage new clinics to participate in this trial. Lately, the Department of Surgery, Kantonsspital Bruderholz, led by PD Dr Igor Langer, has joined in to help us to reach the goal to include 282 patients in the trial. So far over 125 patients have entered the trial.

Furthermore, in collaboration with SAKK project groups we participate as surgeons in other trials, currently in the ongoing lung cancer trials SAKK 16/00 and SAKK 17/04.

Outlook

Prof. Dr Markus Zuber has discontinued his position as vice-chairman of the SAKK Board after two full election periods. All his great achievements are acknowledged. Prof. Walter Marti has been elected as member of the SAKK Board in June 2008 by the General Assembly. This was the moment to renew the presidency of the SAKK Section Surgery. Prof. Paul Schneider, Professor of Surgery, working at the University Hospital of Zurich, being proposed for this position at the annual meeting of the section and elected by the SAKK Board, will take over this position in February 2009. For sure, Prof. Paul Schneider with his vast clinical and research experience will help to boost scientific activities at the Section Surgery.

Section Trial Nurses

President: vacant

Contact person: Christel Böhme, oncology nurse, head of trial coordinators, Oncology/Hematology, Kantonsspital St.Gallen

Short introduction

We are a group of nurses who evaluate draft protocols for their practical and nursing implications and patient considerations, as well as CRF comprehensibility.

In this function we serve as a part of the SAKK internal protocol review process.

Additionally, on each SAKK protocol with a medical treatment a trial nurse is assigned as a contact person for nursing issues.

The group members, with a multifaceted nursing background hold different positions in Swiss hospitals. All work with patients treated in SAKK clinical trials.

Our goal is to make a contribution to assure high-quality clinical trial performance.

Activities 2008

- Review of several SAKK protocols
- Meeting at the semi-annual meeting in Basel
- Exchange of knowledge throughout the year

Outlook

- Continue our work within the Internal Review Board
- Acquire new colleagues to maintain and strengthen our group
- Provide support for nursing issues in ongoing SAKK trials

Network for Cancer Predisposition Testing and Counseling (CPTC)



Presidents:

- 1 PD Dr Pierre O. Chappuis, Unité d'oncogénétique et de prévention des cancers, Service d'Oncologie, University Hospital Geneva (HUG)
- 2 Prof. Dr André-Pascal R. Sappino, Unité d'identification des prédispositions génétiques aux cancers, University Hospital Geneva (HUG)

Short introduction

The main goals of the Network for CPTC are:

- to harmonize the clinical practice of counseling and management of at-risk individuals;
- to consolidate the activity of several Swiss reference molecular laboratories for cancer predisposition testing and the network with basic researchers;
- to collect clinical data and mutation screening results of families with inherited cancer predisposing syndromes that are managed in the network counseling centers;
- to participate in trials evaluating the impact of surveillance and risk reduction strategies;
- to inform and educate health professionals and the lay community on predictive oncology.

Activities 2008

There are 17 centers located in 10 cities throughout Switzerland providing genetic counseling and evaluation for cancer predisposition testing according to the Swiss regulation (KVL/OPAS/OPre art. 12, let. v). More than 400 new families have been managed by network oncogenetic centers this year.

Swiss referral guidelines for genetic counseling and evaluation for BRCA1/BRCA2 testing have been established by the network. These guidelines have been prepared to help clinicians identify the situations where a familial aggregation or a syndrome of hereditary breast/ovarian cancer

should be suspected, and an adequate management could be proposed. These guidelines have been approved by the Swiss Society of Medical Genetics, the Swiss Society of Senology, the Swiss Society of Medical Oncology and the Swiss Society of Gynecology and Obstetrics.

Up to now, complete BRCA1/BRCA2 mutation screening has been performed in 850 distinct families. Members of the network have provided counseling to these families outlining features of the hereditary breast/ovarian cancer syndrome or familial aggregation of breast cancer. Pathogenic mutations have been identified in 20 % of these families and unclassified variants were characterized in 13 % of the index cases tested. We collected clinical and socio-demographic data on families screened for BRCA1/BRCA2 modifications for future research projects.

The IBIS II-Prevention randomized double blind control trial is designed to evaluate anastrozole vs. placebo as an effective method of preventing breast cancer in postmenopausal women at increased risk of the disease. The IBIS II-DCIS trial compares anastrozole vs. tamoxifen as an effective preventive breast cancer drug for postmenopausal women with conservatively-treated ductal in situ cancer. Thirty-two women have already been included in these preventive trials in St.Gallen, Bern, Geneva, Ticino and Lausanne.

Outlook

- to manage individuals identified at high-cancer risk according to standard clinical practice in Switzerland;
- to publish the Swiss guidelines for genetic counseling and evaluation for BRCA1/BRCA2 testing;
- to evaluate the prediction models of germ-line mutations based on the series of families already screened for BRCA1/BRCA2 mutations;
- to initiate a collaborative group to establish guidelines in genetic counseling and testing for patients with multiple colorectal polyps.

Network for Outcomes Research



President:

- 1 Prof. Dr Bernhard Pestalozzi, Department of Oncology, University Hospital Zurich

Vice-President:

- 2 Prof. Dr Thomas Szucs, Institute of Social and Preventive Medicine, University of Zurich

Objectives of the Network

To establish a network of experts and interested researchers and to further promote interdisciplinary outcomes research in oncology. The network may be called in by any SAKK project or working group to provide advice on outcomes research relevant questions in ongoing or planned trials.

Furthermore, the network will collect ideas on outcomes research which can be implemented in oncology trials in Switzerland. One of the important aims of the network is to anchor outcomes research as a part of the clinical trials performed by SAKK. In addition, the network tries to perform other non-randomized controlled trials related outcomes research studies in accordance with the SAKK mission.

Activities 2008 and Outlook

The primary activity is to perform health-economic evaluations (HEA) alongside the clinical trials to provide an incremental cost-effectiveness ratio for a new treatment in question. Although, Switzerland has no institution like the National Institute for Health and Clinical Excellence (NICE) in the U.K. to evaluate cost-effectiveness of drugs, it becomes more and more important to collect health economic information on newly introduced treatments. Sooner or later, this information will be important for health-care decisions.

The Network for Outcomes Research focuses only on direct medical resource usage and its costs. It takes into consideration the perspective of the statutory health insurance or the health system.

For two ongoing SAKK trials the network has evaluated the resource usage and costs of several different data providers. Data from patient records, insurance companies as well as case report forms were evaluated. The evaluation showed that a lot of data are available, but that it is extremely difficult to obtain and combine them in a useful way. Therefore, we decided to use only data from insurance companies in trials where no changes in case report forms are possible anymore. For future trials only case report forms will be used as the primary source to collect data on resource use. Costs are then calculated according to Tarmed or laboratory lists of the Swiss health-care system.

We try to perform cost-effectiveness analyses in different ways. If possible, we will take patient-level costs as they are and perform analyses in line with state of the art methodology, e. g. by using bootstrapping methods to calculate confidence intervals for incremental cost-effectiveness ratios.

For trials with long overall survival times, where it is impossible to record all the cumulated costs and effects over a very long period, costs and effects will have to be modeled.

Trials

SAKK 16/00 (open) | *Preoperative chemoradiotherapy vs. chemotherapy alone in non-small cell lung cancer (NSCLC) patients with mediastinal lymph node metastases (stage IIIA, N2). A randomized prospective phase III trial*

This trial is ongoing and will be considered under a statutory health-insurance perspective, cost data coming from the insurance companies.

SAKK 35/03 (closed) | *Comparing two schedules of rituximab maintenance in rituximab-responding patients with untreated, chemotherapy resistant or relapsed follicular lymphoma. A randomized phase III trial*

This trial has a long-time overall survival and therefore a two-step economic analysis is planned by partially using insurance claim data and modeling cost and effects for long-term survival.

Health-economic analyses for some recently developed trials are planned. For these we will prospectively implement necessary health economic variables into the case report forms to avoid having to collect data from external providers. Although, it is clear that within these forms one cannot

report all resources used, at least the most cost-driving factors such as hospitalizations or expensive treatments can be reported.

We especially want to draw your attention to the trial SAKK 77/08 and SASL 29 *Sorafenib alone or in combination with everolimus in patients with unresectable hepatocellular carcinoma. A randomized multicenter phase II trial*. This trial will be the first in which the resource usage data collection for the HEA, will be completely implemented within the case report forms. Due to the expected short duration of this trial, it serves well as «pilot project» performing HEA this way.

The network is not only involved in SAKK trials but also aims at performing studies in the field of outcomes research in cooperation with third parties, for example studies which are literature-based. Last year, the network performed together with Konstantin Dedes, University of Zurich, a health economic modeling study for the treatment of breast cancer with bevacizumab. This study has been published in the European Journal of Cancer. Similar studies are planned.

Another project in cooperation with the health insurance company Helsana will be a study of end-of-life care in cancer patients in Switzerland. So far, no such study has been performed and therefore, it is unique and has an explorative character. The primary objectives of the study are to evaluate the therapy intensity and hospitalization rates of end-of-life care. Therapy intensity may be defined differently according to literature but we will primarily focus on the percentage of patients (all or sub-groups) receiving any chemo- and/or radiotherapy and/or cancer related surgery during the last month before death. The secondary objective is to assess the magnitude and significance of the effects of insurance coverage, demographic, geographic or treating hospital-related items on the delivery of care.

Depending on the results of this study we hope to initiate a discussion among Swiss oncologists and other cancer treating physicians to reflect on the treatment situation of Swiss cancer patients.

The study will include all identified deceased cancer patients of the years 2006–2008 from the Helsana cohort, but depending on the available number of patients, subgroup analyses for certain cancer types such as lung, colon, breast or prostate are planned.

At the semi-annual SAKK meeting in November 2008 Prof. James Raftery, Director of NETSCC and The Wessex Institute, gave a very interesting talk on «Evidence based decision making: NICE, oncology and health economics.»

Portfolio Plan

SAKK 77/08 and SASL 29 | *Sorafenib alone or in combination with everolimus in patients with unresectable hepatocellular carcinoma. A randomized multicenter phase II trial*

SAKK 77/09 and SASL 30 | *A phase I open label/phase II randomized, double-blind, multicenter trial investigating the combination of everolimus and TransArterial ChemoEmbolisation (TACE) with doxorubicin in patients with hepatocellular carcinoma eligible for TACE. HEA in Phase II*

SAKK 75/08 | *Multimodal therapy with and without cetuximab in patients with locally advanced esophageal carcinoma. An unblinded, prospectively randomized phase III trial*

Collaboration with/Participation in Other Groups

The network initiates project-level cooperation with different institutes active in the field of cancer, e. g. with insurance companies, National Institute for Cancer Epidemiology and Registration (NICER), and the Children's Cancer Registry.

SAKK and Collaborating Groups

Urogenital Tumors

- SAKK 08/91, 08/93, 08/95, 08/97, 08/00** Munger-Beyeler, C., Bernhard, J., Rufibach, K., Morant, R., and Schmid, H.P. (2008). *Quality of analgesic treatment in patients with advanced prostate cancer: do we do a better job now? The Swiss Group for Clinical Cancer Research (SAKK) experience.* Support Care Cancer, 16(5), 461–467. (Journal Impact Factor 2.21)
- EORTC 30891** Studer, U.E., Collette, L., Whelan, P., Albrecht, W., Casselman, J., de Reijke, T., Knonagel, H., Loidl, W., Isorna, S., Sundaram, S.K., and Debois, M. (2008). *Using PSA to guide timing of androgen deprivation in patients with T0-4 N0-2 M0 prostate cancer not suitable for local curative treatment (EORTC 30891).* Eur Urol, 53(5), 941–949. (Journal Impact Factor 5.63)

Lung Cancer

- SAKK 17/00** Ribi, K., Bernhard, J., Schuller, J.C., Weder, W., Bodis, S., Jorger, M., Betticher, D., Schmid, R.A., Stupp, R., Ris, H.B., and Stahel, R.A. (2008). *Individual versus standard quality of life assessment in a phase II clinical trial in mesothelioma patients: feasibility and responsiveness to clinical changes.* Lung Cancer, 61(3), 398–404. (Journal Impact Factor 3.46)
- SAKK 19/03** D'Addario, G., Rauch, D., Stupp, R., Pless, M., Stahel, R., Mach, N., Jost, L., Widmer, L., Tapia, C., Bihl, M., Mayer, M., Ribi, K., Lerch, S., Bubendorf, L., and Betticher, D.C. (2008). *Multicenter phase II trial of gefitinib first-line therapy followed by chemotherapy in advanced non-small-cell lung cancer (NSCLC): SAKK protocol 19/03.* Ann Oncol, 19(4), 739–745. (Journal Impact Factor 4.88)
- EORTC 08971** SILVA Bottomley, A., Debruyne, C., Felip, E., Millward, M., Thiberville, L., D'Addario, G., Rome, L., Zatloukal, P., Coens, C., and Giaccone, G. (2008). *Symptom and quality of life results of an international randomised phase III study of adjuvant vaccination with Bec2/BCG in responding patients with limited disease small-cell lung cancer.* Eur J Cancer, 44(15), 2178–2184. (Journal Impact Factor 4.45)

Breast Cancer

- SAKK 24/06** Greuter S, Schmid F, Ruhstaller T, Thuerlimann B. *Bevacizumab-associated osteonecrosis of the jaw.* Ann Oncol. Letter to the editor. 2008 Dec;19(12):2091–2. Epub 2008 Oct 31. (Journal Impact Factor 4.88)
- IBCSG** Bernhard, J., Zahrieh, D., Zhang, J.J., Martinelli, G., Bassler, R., Hurny, C., Forbes, J.F., Aebi, S., Yeo, W., Thurlimann, B., Green, M.D., Colleoni, M., Gelber, R.D., Castiglione-Gertsch, M., Price, K.N., Goldhirsch, A., and Coates, A.S. (2008). *Quality of life and quality-adjusted survival (Q-TWIST) in patients receiving dose-intensive or standard dose chemotherapy for high-risk primary breast cancer.* Br J Cancer, 98(1), 25–33. (Journal Impact Factor 4.64)
- IBCSG VI** Gruber, G., Cole, B.F., Castiglione-Gertsch, M., Holmberg, S.B., Lindtner, J., Golouh, R., Collins, J., Crivellari, D., Thurlimann, B., Simoncini, E., Fey, M.F., Gelber, R.D., Coates, A.S., Price, K.N., Goldhirsch, A., Viale, G., and Gusterson, B.A. (2008). *Extracapsular tumor spread and the risk of local, axillary and supraclavicular recurrence in node-positive, premenopausal patients with breast cancer.* Ann Oncol, 19(8), 1393–1401. (Journal Impact Factor 4.88)
- IBCSG VII, 12-93** Pagani, O., Gelber, S., Simoncini, E., Castiglione-Gertsch, M., Price, K.N., Gelber, R.D., Holmberg, S.B., Crivellari, D., Collins, J., Lindtner, J., Thurlimann, B., Fey, M.F., Murray, E., Forbes, J.F., Coates, A.S., and Goldhirsch, A. (2008). *Is adjuvant chemotherapy of benefit for postmenopausal women who receive endocrine treatment for highly endocrine-responsive, node-positive breast cancer? International Breast Cancer Study Group Trials VII and 12–93.* Breast Cancer Res Treat. (Journal Impact Factor 4.45)
- IBCSG-VIII** Pestalozzi, B.C., Zahrieh, D., Mallon, E., Gusterson, B.A., Price, K.N., Gelber, R.D., Holmberg, S.B., Lindtner, J., Snyder, R., Thurlimann, B., Murray, E., Viale, G., Castiglione-Gertsch, M., Coates, A.S., and Goldhirsch, A. (2008). *Distinct clinical and prognostic features of infiltrating lobular carcinoma of the breast: combined results of 15 International Breast Cancer Study Group clinical trials.* J Clin Oncol, 26(18), 3006–3014. (Journal Impact Factor 15.48)
- BIG 02-98** Pestalozzi, B.C., Francis, P., Quinaux, E., Dolci, S., Azambuja, E., Gelber, R.D., Viale, G., Baili, A., Andersson, M., Nordenskjold, B., Gnant, M., Gutierrez, J., Lang, I., Crown, J.P., and Piccart-Gebhart, M. (2008). *Is risk of central nervous system (CNS) relapse related to adjuvant taxane treatment in node-positive breast cancer? Results of the CNS substudy in the intergroup Phase III BIG 02-98 Trial.* Ann Oncol, 19(11), 1837–1841. (Journal Impact Factor 4.88)
- BIG 1-98** Viale, G., Giobbie-Hurder, A., Regan, M.M., Coates, A.S., Mastropasqua, M.G., Dell'Orto, P., Maiorano, E., MacGrogan, G., Braye, S.G., Ohlschlegel, C., Neven, P., Orosz, Z., Olszewski, W.P., Knox, F., Thurlimann, B., Price, K.N., Castiglione-Gertsch, M., Gelber, R.D., Gusterson, B.A., and Goldhirsch, A. (2008). *Prognostic and predictive value of centrally reviewed Ki-67 labeling index in postmenopausal women with endocrine-responsive breast cancer: results from Breast International Group Trial 1-98 comparing adjuvant tamoxifen with letrozole.* J Clin Oncol, 26(34), 5569–5575. (Journal Impact Factor 15.48)

BIG 1-02/IBCSG 27-02/NSABP B-37 Wapnir, I. L., Aebi, S., Gelber, S., Anderson, S. J., Lang, I., Robidoux, A., Mamounas, E. P., and Wolmark, N. (2008). *Progress on BIG 1-02/IBCSG 27-02/NSABP B-37, a prospective randomized trial evaluating chemotherapy after local therapy for isolated locoregional recurrences of breast cancer*. *Ann Surg Oncol*, 15(11), 3227–3231. (Journal Impact Factor 3.92)

BIG 1-02/IBCSG 27-02/NSABP B-37 Wapnir, I. L., Aebi, S., Geyer, C. E., Zahrieh, D., Gelber, R. D., Anderson, S. J., Robidoux, A., Bernhard, J., Maibach, R., Castiglione-Gertsch, M., Coates, A. S., Piccart, M. J., Clemons, M. J., Costantino, J. P., and Wolmark, N. (2008). *A randomized clinical trial of adjuvant chemotherapy for radically resected locoregional relapse of breast cancer: IBCSG 27-02, BIG 1-02, and NSABP B-37*. *Clin Breast Cancer*, 8(3), 287–292.

Leukemia

SAKK 31/98 Zenhausem, R., Simcock, M., Gratwohl, A., Hess, U., Bargetzi, M., and Tobler, A. (2008). *Rituximab in patients with hairy cell leukemia relapsing after treatment with 2-chlorodeoxyadenosine (SAKK 31/98)*. *Haematologica*, 93(9), 1426–1428. (Journal Impact Factor 5.52)

APL Ades, L., Sanz, M. A., Chevret, S., Montesinos, P., Chevalier, P., Raffoux, E., Vellenga, E., Guerci, A., Pignieux, A., Huguet, F., Rayon, C., Stoppa, A. M., de la Serna, J., Cahn, J.-Y., Meyer-Monard, S., Pabst, T., Thomas, X., de Botton, S., Parody, R., Bergua, J., Lamy, T., Vekhoff, A., Negri, S., Ifrah, N., Dombret, H., Ferrant, A., Bron, D., Degos, L., and Fenau, P. (2008). *Treatment of newly diagnosed acute promyelocytic leukemia (APL): a comparison of French-Belgian-Swiss and PETHEMA results*. *Blood*, 111, 1078–84. (Journal Impact Factor 10.90)

Lymphoma

IFM99-03 and IFM99-04 Moreau, P., Garban, F., Attal, M., Michallet, M., Marit, G., Hulin, C., Benboubker, L., Doyen, C., Mohty, M., Yakoub-Agha, I., Leyvraz, S., Casassus, P., Avet-Loiseau, H., Garderet, L., Mathiot, C., and Harousseau, J. L. (2008). *Long-term follow-up results of IFM99-03 and IFM99-04 trials comparing nonmyeloablative allotransplantation with autologous transplantation in high-risk de novo multiple myeloma*. *Blood*, 112(9), 3914–3915. (Journal Impact Factor 10.90)

DSHNHL Klapper W, Stoecklein H, Zeynalova S, Ott G, Kosari F, Rosenwald A, Loeffler M, Trümper L, Pfreundschuh M, Siebert R; German High-Grade Non-Hodgkin's Lymphoma Study Group. *Structural aberrations affecting the MYC locus indicate a poor prognosis independent of clinical risk factors in diffuse large B-cell lymphomas treated within randomized trials of the German High-Grade Non-Hodg-*

kin's Lymphoma Study Group (DSHNHL). *Leukemia*. 2008 Dec;22(12):2226–9. Epub 2008 Aug 28. (Journal Impact Factor 6.92)

GHSg Sieniawski, M., Reineke, T., Josting, A., Nogova, L., Behringer, K., Halbsguth, T., Fuchs, M., Diehl, V., and Engert, A. (2008). *Assessment of male fertility in patients with Hodgkin's lymphoma treated in the German Hodgkin Study Group (GHSg) clinical trials*. *Ann Oncol*, 19(10), 1795–1801. (Journal Impact Factor 4.88)

GHSg-HD15 Kobe, C., Dietlein, M., Franklin, J., Markova, J., Lohri, A., Amthauer, H., Klutmann, S., Knapp, W. H., Zijlstra, J. M., Bockisch, A., Weckesser, M., Lorenz, R., Schreckenberger, M., Bares, R., Eich, H. T., Mueller, R. P., Fuchs, M., Borchmann, P., Schicha, H., Diehl, V., and Engert, A. (2008). *Positron emission tomography has a high negative predictive value for progression or early relapse for patients with residual disease after first-line chemotherapy in advanced-stage Hodgkin lymphoma*. *Blood*, 112(10), 3989–3994. (Journal Impact Factor 10.90)

MINT Pfreundschuh M, Ho AD, Cavallin-Stahl E, Wolf M, Pettengell R, Vasova I, Belch A, Walewski J, Zinzani PL, Mingrone W, Kvaloy S, Shpilberg O, Jaeger U, Hansen M, Corrado C, Scheliga A, Loeffler M, Kuhnt E; MabThera International Trial (MINT) Group. (2008) *Prognostic significance of maximum tumour (bulk) diameter in young patients with good-prognosis diffuse large-B-cell lymphoma treated with CHOP-like chemotherapy with or without rituximab: an exploratory analysis of the MabThera International Trial Group (MINT) study*. *Lancet Oncol*;9:435–44. (Journal Impact Factor 12.25)

Gastrointestinal Cancer

SAKK 40/87 Laffer, U., Metzger, U., Aeberhard, P., Lorenz, M., Harder, F., Maibach, R., Zuber, M., and Herrmann, R. (2008). *Adjuvant perioperative portal vein or peripheral intravenous chemotherapy for potentially curative colorectal cancer: long-term results of a randomized controlled trial*. *Int J Colorectal Dis*, 23(12), 1233–1241. (Journal Impact Factor 1.92)

SAKK 41/04 Borner, M., Koeberle, D., Von Moos, R., Saletti, P., Rauch, D., Hess, V., Trojan, A., Helbling, D., Pestalozzi, B., Caspar, C., Ruhstaller, T., Roth, A., Kappeler, A., Dietrich, D., Lanz, D., and Mingrone, W. (2008). *Adding cetuximab to capecitabine plus oxaliplatin (XELOX) in first-line treatment of metastatic colorectal cancer: a randomized phase II trial of the Swiss Group for Clinical Cancer Research SAKK*. *Ann Oncol*, 19(7), 1288–1292. (Journal Impact Factor 4.88)

SAKK 44/00 Bernhard, J., Dietrich, D., Scheithauer, W., Gerber, D., Bodoky, G., Ruhstaller, T., Glimelius, B., Bajetta, E., Schuller, J., Saletti, P., Bauer, J., Figer, A., Pestalozzi, B. C., Kohne, C. H., Mingrone, W., Stemmer, S. M., Tamas, K.,

Kornek, G. V., Koeberle, D., and Herrmann, R. (2008). *Clinical benefit and quality of life in patients with advanced pancreatic cancer receiving gemcitabine plus capecitabine versus gemcitabine alone: a randomized multicenter phase III clinical trial-SAKK 44/00-CECOG/PAN. 1.3.001*. *J Clin Oncol*, 26(22), 3695–3701. (Journal Impact Factor 15.48)

SAKK 44/00 Hess V, Glimelius B, Grawe P, Dietrich D, Bodoky G, Ruhstaller T, Bajetta E, Saletti P, Figer A, Scheithauer W, Herrmann R. *CA 19-9 tumour-marker response to chemotherapy in patients with advanced pancreatic cancer enrolled in a randomised controlled trial*. *Lancet Oncol*. 2008 Feb;9(2):132–8. (Journal Impact Factor 12.25)

SAKK 44/02 Koeberle, D., Saletti, P., Borner, M., Gerber, D., Dietrich, D., Caspar, C. B., Mingrone, W., Beretta, K., Strasser, F., Ruhstaller, T., Mora, O., and Herrmann, R. (2008). *Patient-reported outcomes of patients with advanced biliary tract cancers receiving gemcitabine plus capecitabine: a multicenter, phase II trial of the Swiss Group for Clinical Cancer Research*. *J Clin Oncol*, 26(22), 3702–3708. (Journal Impact Factor 15.48)

SAKK 75/06 single case Grimm B, Padberg B, Ruhstaller T, Fleisch F, von Moos R. *Fatal herpes simplex virus hepatitis in a patient with esophageal cancer under radiochemotherapy*. *Onkologie*. 2008 Nov;31(11):620-2. Epub 2008 Oct 27. (Journal Impact Factor 1.23)

Melanoma

EORTC 18991 Eggermont, A. M., Suci, S., Santinami, M., Testori, A., Kruit, W. H., Marsden, J., Punt, C. J., Sales, F., Gore, M., Mackie, R., Kusic, Z., Dummer, R., Hauschild, A., Musat, E., Spatz, A., and Keilholz, U. (2008). *Adjuvant therapy with pegylated interferon alfa-2b versus observation alone in resected stage III melanoma: final results of EORTC 18991, a randomised phase III trial*. *Lancet*, 372(9633), 117–126. (Journal Impact Factor 28.64)

Sarcoma Pieper S, Ranft A, Braun-Munzinger G, Jurgens H, Paulussen M, Dirksen U. *Ewing's tumors over the age of 40: a retrospective analysis of 47 patients treated according to the International Clinical Trials EICESS 92 and EURO-E. W. I. N. G. 99*. *Onkologie*. 2008 Dec;31(12):657–63. Epub 2008 Nov 20. (Journal Impact Factor 1.23)

Central Nervous System Tumors

EORTC 26981 Preusser, M., Charles Janzer, R., Felsberg, J., Reifenberger, G., Hamou, M. F., Diserens, A. C., Stupp, R., Gorlia, T., Marosi, C., Heinzl, H., Hainfellner, J. A., and Hegi, M. (2008). *Anti-O6-methylguanine-methyltransferase (MGMT) immunohistochemistry in glioblastoma multiforme: observer variability and lack of association with patient survival impede its use as clinical biomarker*. *Brain Pathol*, 18(4), 520–532. (Journal Impact Factor 3.66)

Lamers, L. M., Stupp, R., van den Bent, M. J., Al, M. J., Gorlia, T., Wasserfallen, J. B., Mittmann, N., Jin Seung, S., Crott, R., and Uyl-de Groot, C. A. (2008). *Cost-effectiveness of temozolomide for the treatment of newly diagnosed glioblastoma multiforme: a report from the EORTC 26981/22981 NCI-C CE3 Intergroup Study*. *Cancer*, 112(6), 1337–1344. (Journal Impact Factor 4.63)

Gorlia T, van den Bent MJ, Hegi ME, Mirimanoff RO, Weller M, Cairncross JG, Eisenhauer E, Belanger K, Brandes AA, Allgeier A, Lacombe D, Stupp R. *Nomograms for predicting survival of patients with newly diagnosed glioblastoma: prognostic factor analysis of EORTC and NCIC trial 26981-22981/CE.3*. *Lancet Oncol* 9:29–38, 2008. (Journal Impact Factor 12.25)

Stupp R, Hottinger AF, van den Bent MJ, Dietrich PY, Brandes AA. *Frequently asked questions in the medical management of high-grade glioma: a short guide with practical answers*. *Ann Oncol* 19 (Suppl 7):vii209–16, 2008. (Journal Impact Factor 4.88)

Wick W, Stupp R, Beule AC, Bromberg J, Wick A, Ernemann U, Platten M, Marosi C, Mason WP, van den Bent M, Weller M, Rorden C, Karnath HO; The European Organisation for Research and Treatment of Cancer and the National Cancer Institute of Canada Clinical Trials Group. *A novel tool to analyze MRI recurrence patterns in glioblastoma*. *Neuro Oncol*. 2008 Aug 4. [Epub ahead of print] (Journal Impact Factor 5.80)

Murat A, Migliavacca E, Gorlia T, Lambiv WL, Shay T, Hamou MF, de Tribolet N, Regli L, Wick W, Kouwenhoven MCM, Hainfellner JA, Heppner FL, Dietrich PY, Zimmer Y, Cairncross JG, Janzer RC, Domany E, Delorenzi M, Stupp R, and Hegi ME. *Stem Cell-Related «Self-renewal signature» and high Epidermal Growth Factor Receptor expression associated with resistance to concomitant chemo-radiotherapy in glioblastoma*. *J Clin Oncol* 26: 3015–3024, 2008. (Journal Impact Factor 15.48)

Palliative Cancer Care

Strasser, F., Demmer, R., Bohme, C., Schmitz, S. F., Thuerlimann, B., Cerny, T., and Gillessen, S. (2008). *Prevention of docetaxel- or paclitaxel-associated taste alterations in cancer patients with oral glutamine: a randomized, placebo-controlled, double-blind study*. *Oncologist*, 13(3), 337–346. (Journal Impact Factor 4.88)

Strasser F, Lutz TA, Maeder MT, Thuerlimann B, Bueche D, Tschöp M, Kaufmann K, Holst B, Brändle M, von Moos R, Demmer R, Cerny T. *Safety, tolerability and pharmacokinetics of intravenous ghrelin for cancer-related anorexia/cachexia: a randomised, placebo-controlled, double-blind, double-crossover study.* Br J Cancer. 2008 Jan 29;98(2):300–8. Epub 2008 Jan 8. (Journal Impact Factor 4.64)

Outcomes Research

Matter-Walstra, K., Szucs, T., Brauchli, P., Schwenkglenks, M. (2008). *Randomisiert-kontrollierte Studien von Krebstherapien in der Schweiz. Outcomes Research und gesundheitsökonomische Aspekte.* Schweizerische Ärztezeitung, 2008;89(36).

Cancer Predisposition Testing and Counseling

CAPP2 Burn J, Bishop DT, Mecklin JP, Macrae F, Möslin G, Olschwang S, Bisgaard ML, Ramesar R, Eccles D, Maher ER, Bertario L, Jarvinen HJ, Lindblom A, Evans DG, Lubinski J, Morrison PJ, Ho JW, Vasen HF, Side L, Thomas HJ, Scott RJ, Dunlop M, Barker G, Elliott F, Jass JR, Fodde R, Lynch HT, Mathers JC. *Effect of aspirin or resistant starch on colorectal neoplasia in the Lynch syndrome.* N Engl J Med. 2008 Dec 11;359(24):2567–78. (Journal Impact Factor 52.59)

Liljegren, A., Barker, G., Elliott, F., Bertario, L., Bisgaard, M.L., Eccles, D., Evans, G., Macrae, F., Maher, E., Lindblom, A., Rotstein, S., Nilsson, B., Mecklin, J.P., Moslein, G., Jass, J., Fodde, R., Mathers, J., Burn, J., and Bishop, D.T. (2008). *Prevalence of adenomas and hyperplastic polyps in mismatch repair mutation carriers among CAPP2 participants: report by the colorectal adenoma/carcinoma prevention programme 2.* J Clin Oncol, 26(20), 3434–3439. (Journal Impact Factor 15.48)

Kote-Jarai Z, Easton DF, Stanford JL, Ostrander EA, Schleutker J, Ingles SA, Schaid D, Thibodeau S, Dörk T, Neal D, Donovan J, Hamdy F, Cox A, Maier C, Vogel W, Guy M, Muir K, Lophatananon A, Kedda MA, Spurdle A, Steginga S, John EM, Giles G, Hopper J, Chappuis PO, Hutter P, Foulkes WD, Hamel N, Salinas CA, Koopmeiners JS, Karyadi DM, Johanneson B, Wahlfors T, Tammela TL, Stern MC, Corral R, McDonnell SK, Schürmann P, Meyer A, Kuefer R, Leongamornlert DA, Tymrakiewicz M, Liu JF, O'Mara T, Gardiner RA, Aitken J, Joshi AD, Severi G, English DR, Southey M, Edwards SM, Al Olama AA; PRACTICAL Consortium, Eeles RA. *Multiple novel prostate cancer predisposition loci confirmed by an international study: the PRACTICAL Consortium.* Cancer Epidemiol Biomarkers Prev. 2008 Aug;17(8):2052-61. Erratum in: Cancer Epidemiol Biomarkers Prev. 2008 Oct;17(10):2901. Donovan, Jenny [added]; Hamdy, Freddie [added]. (Journal Impact Factor 4.64)

Linger, R., Dudakia, D., Huddart, R., Tucker, K., Friedlander, M., Phillips, K. A., Hogg, D., Jewett, M. A., Lohynska, R., Daugaard, G., Richard, S., Chompret, A., Stoppa-Lyonnet, D., Bonaiti-Pellie, C., Heidenreich, A., Albers, P., Olah, E., Geczi, L., Bodrogi, I., Daly, P. A., Guilford, P., Fossa, S. D., Heimdal, K., Tjulandin, S. A., Liubchenko, L., Stoll, H., Weber, W., Einhorn, L., McMaster, M., Korde, L., Greene, M. H., Nathanson, K. L., Cortessis, V., Easton, D. F., Bishop, D. T., Stratton, M. R., and Rapley, E. A. (2008). *Analysis of the DND1 gene in men with sporadic and familial testicular germ cell tumors.* Genes Chromosomes Cancer, 47(3), 247–252. (Journal Impact Factor 4.53)

Novak, D. J., Sabbaghian, N., Maillet, P., Chappuis, P. O., Foulkes, W. D., and Tischkowitz, M. (2008). *Analysis of the genes coding for the BRCA1-interacting proteins, RAP80 and Abraxas (CCDC98), in high-risk, non-BRCA1/2, multi-ethnic breast cancer cases.* Breast Cancer Res Treat. (Journal Impact Factor 4.45)

Roh V, Laemmle A, Von Holzen U, Stroka D, Dufour JF, Hunt KK, Candinas D, Vorburger SA. *Dual induction of PKR with E2F-1 and IFN-alpha to enhance gene therapy against hepatocellular carcinoma.* Cancer Gene Ther. 2008 Oct;15(10):636–44. Epub 2008 Jun 6. (Journal Impact Factor 3.89)

Resoconto annuale dal 1° gennaio al 31 dicembre (in CHF)	2008	2007
Reddito di esercizio		
Contributi per la ricerca SBF ¹	3 930 520.00	5 266 800.00
Altri contributi per la ricerca ²	1 427 794.58	765 860.89
Ricavi collaborazioni industriali	2 268 112.60	2 319 453.15
Redditi bollettino sul cancro	281 885.00	352 655.00
Donazione, Legati & Eredità	40 720.00	0.00
Redditi vari	484 918.85	211 564.60
Totale reddito di esercizio	8 433 951.03	8 916 333.64
Spese di esercizio		
Costi per studi vari	-529 278.75	-311 799.90
Contributi per la ricerca SPOG ³	0.00	-449 460.00
Contributi per la ricerca ASRT (NICER) ⁴	0.00	-900 900.00
Contributi per la ricerca IBCSG ⁵	-250 000.00	-250 000.00
Contributi per la ricerca Centri	-2 699 123.95	-2 198 918.25
Costi di trasferta e viaggi di rappresentanza	-241 480.75	-199 901.50
Altre spese di esercizio	-63 649.62	-154 901.55
Totale spese di esercizio	-3 783 533.07	-4 465 881.20
Subtotale 1	4 650 417.96	4 450 452.44
Costi di coordinamento		
Costi per il personale	-4 080 476.39	-3 160 505.11
Altri costi di coordinamento	-831 077.70	-975 575.25
Totale costi di coordinamento	-4 911 554.09	-4 136 080.36
Subtotale 2	-261 136.13	314 372.08
Risultato finanziario		
Reddito finanziario	131 068.35	50 618.64
Oneri finanziari	-117 838.11	-2 194.80
Totale risultato finanziario	13 230.24	48 423.84
Subtotale 3	-247 905.89	362 795.92
Attività al di fuori del periodo		
Attività al di fuori del periodo considerato	58 024.39	0.00
Scioglimento di accantonamenti non necessari	154 628.40	77 900.00
Total résultat exceptionnel et hors exploitation	212 652.79	77 900.00
Totale attività al di fuori del periodo considerato	-35 253.10	440 695.92

1 2007: Segreteria di stato per l'educazione e la ricerca SER (mediante il SIAK i fondi vanno alle relative associazioni, come de accordo sulle prestazioni con l'unione) A partire dal 2008 i fondi sono passati direttamente dalla SER al SAKK, allo SPOG e alla ASRT (NICER)

2 Lega contro il cancro/Ricerca svizzera contro il cancro/Fondazione svizzera per la ricerca clinica contro il cancro/Oncosuisse

3 Gruppo Onco-Pediatico Svizzero SPOG

4 Associazione Svizzera Registri Tumori ASRT

5 International Breast Cancer Study Group IBCSG

Schweizerische Arbeitsgemeinschaft für Klinische Krebsforschung SAKK
Groupe Suisse de Recherche Clinique sur le Cancer SAKK
Swiss Group for Clinical Cancer Research SAKK
Gruppo Svizzero di Ricerca Clinica sul Cancro SAKK

SAKK Coordinating Center

Effingerstrasse 40
3008 Berna
Tel. 031 389 91 91
Fax 031 389 92 00
<http://sakk.ch>
sakkcc@sakk.ch

Relazione bancaria (per offerte)

Banca Cantonale di Berna
BEKB, Bundesplatz 8
3001 Berna
IBAN CH14 0079 0042 4106 0712 5

SAKK Industriepool 2008

**Ringraziamo I gruppi farmaceutici
per il loro sostegno:**

Amgen Switzerland AG
AstraZeneca AG
Baxter AG
Bayer Schering Pharma (Schweiz) AG
Bristol-Myers Squibb GmbH
Celgene GmbH
Eli Lilly Suisse S. A.
Essex Chemie AG
GlaxoSmithKline AG
Janssen-Cilag AG
Lipomed AG
Mepha Pharma AG
Merck (Schweiz) AG
Mundipharma Medical Company
Novartis Pharma Schweiz AG
Nycomed Pharma AG
Pfizer AG
Robapharm AG
Roche Pharma Schweiz AG
sanofi-aventis (suisse) sa
Wyeth Pharmaceuticals AG

Contributi dal settore pubblico e d'altri

Segreteria di Stato per l'educazione e la ricerca SER
Ricerca Svizzera contro il Cancro
Oncosuisse
Lega contro il cancro Svizzera
Schweizerische Stiftung für Klinische Krebsforschung
(Fondazione svizzera per la ricerca clinica contro il cancro)
Donatori privati

Redazione

Ruth Kellenberger

Traduzione

Mirta Piller, Zurigo

Realizzazione

atelierrichner.ch

Stampa

RMS Repro Media Services AG, Berna

Immagini

Peter Schneider, Thun

SAKK Coordinating Center

Effingerstrasse 40

3008 Berna

Tel. 031 389 91 91

Fax 031 389 92 00

<http://sakk.ch>

sakkcc@sakk.ch

